

Determine the frequency of Bacterial Pathogens in Neonatal Sepsis and their Sensitivity to Different Antibiotics.

Muhammad Sohail¹, Mohsina Haq^{2*}, Bakht Biland Khan³, Shah Zaman⁴, Amir Zaman Khan⁴, Saima mehboob⁵, Ihteshamul Haq^{6*}, Jauhar Nafees⁷, Alia Gul⁸, Muhammad Umair Khan⁹, Obaid Ullah Shah¹⁰,

1. Department of Medical Laboratory Technology, Faculty of Rehabilitation & Allied Health Sciences, Riphah International University Islamabad, QIE Campus, Lahore, Pakistan
2. Department of Pathology Peshawar Medical college, Riphah international University, Warsak road Peshawar Kp Pakistan
3. Department of Medicine, Mercy Teaching Hospital, Peshawar Medical College Kp Pakistan
4. Department of Pharmacology, Peshawar Medical College Warsak road Peshawar Kp Pakistan
5. Department of Medicine Kuwait teaching hospital, Peshawar Medical college Warsak road Peshawar Kp Pakistan
6. Department of Biotechnology and Genetic Engineering Hazara University Mansehra Kp Pakistan
7. Department of Zoology Government Degree College Lund Khwar Mardan, Abdul Wali Khan University Mardan Kp Pakistan
8. Department of Botany Hazara University Mansehra Kp Pakistan
9. Department of Biotechnology Abdul Wali Khan University Mardan kp Pakistan
10. College of Tropical Crops, Hainan University, 570228 PR China.

Corresponding Author: Mohsina Haq^{2*}

Email Address: dr.mohsinahaq@gmail.com

Co- Corresponding Author: Ihteshamul Haq^{6*}

Email Address: ihteshamulhaq384@gmail.com

Abstract:

Neonatal sepsis is a blood infection that occurs in an infant younger than 90 days old. Early onset sepsis is seen in the first week of life. Late onset sepsis occurs after 1 week through 3 months of

age. The aim of the present research to evaluate the Prevalence rate of Bacterial pathogens in neonatal sepsis and to determine the sensitivity of these pathogens to commonly used antibiotics at Department of Pediatrics, at PIMS Hospital Islamabad Pakistan from January 2016 to December 2016. In the current study 100 neonates were enrolled and Sample for Blood culture was taken under aseptic measures by standardized techniques and were saved in culture bottle. It was collected and transported to the Hospital laboratory by single person and inoculated & incubated there. The growth of the pathogen was observed for next 5 days and sensitivity pattern was checked by using different antibiotic discs. Culture and sensitivity pattern as shown by blood culture report was noted. The result show that the Frequency of bacterial pathogens were 35(34%) Klebsiella, 28(26%) Pseudomonas, 22(20%) E.coli and Staphylococcus 26(24%), while frequency of antimicrobial sensitivity reveals 77% Ciprofloxacin, 83% Amikacin, 19% Penicillin and 12% Gentamicin for Staphylococcus Aureus. Sensitivity for Pseudomonas was 58% Ciprofloxacin, 61% Amikacin, 1% Penicillin, Imipenem (81%) and 37% Gentamicin. For pathogens Klebsiella it was 47% Ciprofloxacin, 37% Amikacin, and 9% Imipenem while Ciprofloxacin 49%, Amikacin 54% and Imipenem 78% for E.coli. From The Present Research we concluded that Klebsiella and Pseudomonas are the leading pathogens in neonatal sepsis and the most commonly used antibiotics using in our setup are highly resistant with them which may be used when culture reports are awaiting.

Keywords: Neonatal sepsis, pathogens, antibiotics, resistance

INTRODUCTION

Neonatal sepsis is a blood infection that occurs in an infant younger than 90 days old. Early onset sepsis is seen in the first week of life. Late onset sepsis occurs after 1 week through 3 months of age. Neonatal sepsis may be categorized as early or late onset. Early onset sepsis has onset in the first 3 days of life and late onset sepsis is sepsis occurring during 4-90 days of life (1). The organisms most commonly associated with early-onset infection include Group B Streptococcus (GBS), Escherichia coli, coagulase negative Staphylococcus, Haemophilus influenza, and Listeria monocytogenes (3). These organisms are acquired from the mother during birth. Late-onset sepsis is acquired from the environment. Organisms that have been implicated in causing late-onset sepsis syndrome include coagulase-negative staphylococci, Staphylococcus aureus, E coli, Klebsiella, Pseudomonas, Enterobacter, Candida, GBS, Serratia, Acinetobacter and

anaerobe (4) Incidence of Neonatal sepsis in clinically diagnosed cases is 170/1000 live births, and in cases confirmed by blood culture the incidence is 5.5/1000 live births (5). Neonatal Sepsis is one of the main cause of mortality and morbidity in neonates especially in the developing countries. In Pakistan neonatal mortality rate is 47.3 per 1000 live births; 23% of the deaths being the results of infections (6,7). According to a study conducted in Pakistan, cases of culture proven sepsis were 54%; 42% being early onset sepsis and 58% late onset sepsis (7,8).Percentage of various pathogens in early onset sepsis is Klebsiella species (25%), Escherichia coli (15%), and Staphylococcus aureus (18%), Group B streptococcus (7%) and in late onset sepsis is Staphylococcus aureus (14%), GBS (12%), Streptococcus pneumoniae (12%), and nontyphoidal Salmonella species (13%) (9,10).Neonatal sepsis (NS) may have subtle, diverse and nonspecific symptoms and signs; moreover, a delay in the diagnosis and commencement of treatment results in a high morbidity and mortality⁸. In our county pre-existing data on both early and late onset sepsis has shown great diversity in the changing pattern of the organism and their sensitivity patterns⁹. Most of the bacterial agents are resistant to amoxicillin/ampicillin and third generation cephalosporins (10,11).We designed this study with the vision that the resistance of bacterial pathogens to commonly used antibiotics is emerging and therefore sensitivity of these pathogens to antibiotics is changing. So it was the need of time to reconsider the frequency of bacterial organisms and the recent sensitivity pattern as it is changing due to emerging antibiotic resistance. This data will give guideline for experimental use of antibiotics when culture reports are awaited.

MATERIAL AND METHODS

A total of 100 Neonate (1 to 28 days) both male and females from Department of Pediatrics, at PIMS Hospital Islamabad Pakistan were included in the study and neonates having life threatening co morbid illness i.e. Birth Asphxia (Negative Blood Culture) and Meconium Aspiration Syndrome (Negative Blood Culture) were excluded from the study. Informed consent was taken from the parents. Demographics like name, age, sex and address were noted. Sample for Blood culture was taken under aseptic measures by standardized techniques and were saved in culture bottle. It was collected and transported to the Services Hospital Lahore laboratory by single person and inoculates & incubated there. The growth of the pathogen was observed for next 5 days and sensitivity pattern was checked by using different antibiotic discs. All this was

done in same laboratory so that results are not altered. Culture and sensitivity pattern as shown by blood culture report was noted. Information regarding data of the patient, type of the pathogen in culture and its susceptibility to antibiotics were entered in a Proforma. The data was entered and analyzed by using SPSS 10 computer system. Descriptive Statistics was applied in the form of frequencies and percentages for qualitative variables like gender, bacterial pathogens and their antimicrobial sensitivity. Mean \pm standard deviation for quantitative variables like age of the neonate.

RESULTS

Age distribution of the patients was done, majority of the patients were recorded between 11-20 days i.e., 59(49%), 36(36%) between 1-10 days and 15(15%) between 21-28 days, while mean and sd was calculated as 18.54 \pm 8.76 days. Gender distribution of the patients was done in Table 2, where 43(45%) were male and 57(55%) females. Frequency of bacterial pathogens were recorded 35(34%) Klebsiella, 28(26%) Pseudomonas, 22(20%) E.coli and Staphylococcus 26(24%). Frequency of antimicrobial sensitivity was done for pathogens which reveals 77% Ciprofloxacin, 83% Amikacin, 19% Penicillin and 12% Gentamicin for Staphylococcus Aureus. Sensitivity for Pseudomonas was 58% Ciprofloxacin, 61% Amikacin, 1% Penicillin, Imipenem (81%) and 37% Gentamicin. For pathogens Klebsiella it was 47% Ciprofloxacin, 37% Amikacin, and 9% Imipenem while Ciprofloxacin 49%, Amikacin 54% and Imipenem 78% for E.coli. (Table 1)

Table 1: Frequency of Antimicrobial Sensitivity (n=100)

Pathogens	Sensitivity
Staphylococcus Aureus	Ciprofloxacin (77%), Amikacin(83%), Penicillin(19%), Gentamicin(12%)
Pseudomonas	Ciprofloxacin (58%), Amikacin(61%), Penicillin(1%), Gentamicin(37%) Imipenem (81%)
Klebsiella	Ciprofloxacin (47%), Amikacin(37%), Imipenem(9%)

E.coli	Ciprofloxacin (49%), Amikacin(54%), Imipenem(78%)
--------	---

DISCUSSION

Neonatal sepsis is an important and common cause of neonatal morbidity and mortality. The incidence of neonatal sepsis in the developed countries is 1- 10/1000 live births,(12,13). Where as it is roughly three times more in developing countries like Pakistan.(14,15). This high incidence is mainly due to poor antenatal care and lack of trained staff to conduct deliveries.(16,17)There is a strong association between maternal urinary tract infection, pyrexia, vaginal discharge & unclean vaginal examination during labour and early onset neonatal sepsis (18,19).We planned this study to determine the frequency of underlying pathogens in neonatal sepsis and to determine the sensitivity of these pathogens to commonly used antibiotics as the resistance of bacterial pathogens to commonly used antibiotics is emerging and therefore sensitivity of these pathogens to antibiotics is changing. The findings of study are in agreement with a study where Klebsiella species was (25%), Escherichia coli (15%), and Staphylococcus aureus (18%)(20,21). Another study report show that Klebsiella pneumoniae as the predominant pathogen (36.7%), followed by Pseudomonas species (30%) and recorded most of isolates highly resistant to the majority of other antibiotics tested (22,23). However, we revealed that the resistance of bacterial pathogens to commonly used antibiotics are resistant. The results further give us guideline for empirical use of antibiotics when culture reports are awaited.

CONCLUSION

The results of the study show that Klebsiella and Pseudomonas are the leading pathogens in neonatal sepsis and the most commonly used antibiotics using in our setup are highly resistant with them which may be used when culture reports are awaiting.

REFERENCE

1. Ann L Anderson-Berry, MD, Neonatal sepsis emedicine specialities Pediatrics: Cardiac Disease and Critical Care Medicine, Neonatology 2009. www.emedicine.com/article/978352-overview

2. ul Haq, I., Khan, M., Rehman, Z., Anwar, F., Ullah, H., & Ullah, N. (2018). HBV prevalence in the volunteer blood donors in Peshawar, Khyber Pakhtunkhwa Pakistan. *Int J Biosci*, 13(5), 50-54.
3. van den Hoogen A, Gerards LJ, Verboon-Maciolek MA, Fleer A, Krediet TG. Long-Term Trends in the Epidemiology of Neonatal Sepsis and Antibiotic Susceptibility of Causative Agents. *Neonatology*. 2009;97(1):22-8.
4. Thaver, Durrane , Zaidi, Anita K. M. Burden of Neonatal Infections in Developing Countries: A Review of Evidence From Community-Based Studies. *The Pediatric Infectious Disease Journal*. 2009;28(1):suppl S3-S9.
5. Khalid, F. (2021). Comparative Diagnostic Analysis and Biochemical Profile in Patients with Covid-19, Dengue and Acute Febrile Illness: Suggestions for Patient Controlling. *Annals of the Romanian Society for Cell Biology*, 25(7), 1733-1744.
6. Aftab R, Iqbal I. Bacteriological agents of Neonatal Sepsis in NICU at Nishtar Hospital Multan. *JCPSP*. March 2006;16(3):216-9.
7. Rehman, A., Haq, I., Asghar, Afridi, G. Z., & Faisal, S. (2020). Sero-epidemiological Identification of Dengue Virus in Individuals at District Shangla, Khyber Pakhtunkhwa. Pakistan. *J Biomedical Sci*, 9(3),10
8. Ahmed Z, Ghafoor T, Waqar T. Diagnostic Value Of Creactive protein And Haematological Parameters In Neonatal Sepsis. *JCPSP* 2005;15:152-6.
9. Butt-M.A, Malik-BA, Sheikh-S, Shamoan M. Infections in Preterm The Professional medical journal. 2005;12(4);451-6.
10. Kaistha N, Mehta M, Singla N, Garg R, Chander J. Neonatal septicemia isolates and resistance patterns in a tertiary care hospital of North India. *Journal of Infection in developing countries*.2009;13:4(1):55-7.
11. Jamal-M, Khan N. Neonatal Morbidity and Mortality in high risk pregnancies. *JCPSP* 2002;12(11):657-61.
12. Haq, I., Muhammad, A., Fazli Zahir, M. K., Anwar, F., Akhtar, M. S., & Ullah, F. (2020). Serological and Epidemiology study of Helicobacter pylori infection among Dyspeptic patients in District Peshawar Pakistan. *Adv. Biores*, 11(3), 81-85

13. Bhutta ZA, Yusuf K. Early onset neonatal sepsis in Pakistan. A case control study of birth cohort. *Am J Perinatology* 1997;14(a):557-81.
14. Dias E. The bacterial profile of neonatal septicaemia in a rural hospital in south india. *Journal of Clinical and Diagnostic Research*.2010;4(6):3328-31.
15. Al-Shamahy HA, Sabrah AA, Al-Robasi AB, Naser SM. Types of Bacteria associated with Neonatal Sepsis in Al-Thawra University Hospital, Sana'a, Yemen, and their Antimicrobial Profile. *SQU Med J* 2012;12:48-54.
16. Anwar, F., Tayyab, M., Haq, I., & Shah, O. U. (2021). Viral overload of COVID-19 pandemics: Overweight people a soft target to get an infection. *International Journal of Clinical Virology*,5(2),070-071.
17. Qamar, Z., Anwar, F., Ahmad, R., Haq, I., Khan, A. M. K., Hussain, R., ... & Khan, J. (2021). Prevalence of Hepatitis C virus and determination of its genotypes in subjects of Tehsil Daggar District Buner,KP,Pakistan. *Clinical Epidemiology and Global Health*, 12, 100809.
18. Asghar, M., Haq, M., & Saleem, N. (2022). Molecular Identification And Prevalence Rate Of HepatitisC Virus Among Hemodialysis Patients In Peshawar Kp Pakistan. *NVEO-NATURAL VOLATILES & ESSENTIAL OILS Journal| NVEO*, 1168-1175
19. Asif, A., Asghar, M., Khan, H. U., Haq, I., Shuaib, S. L., Khalid, F., ... & Rehman, N. (2021). Antibiotic susceptibility pattern of clinical isolates of methicillin resistant staphylococcus aureus in Peshawar,Pakistan. *Annals of the Romanian Society for Cell Biology*, 25(6), 20116-20131
20. Haq, I., Zahir, F., ur Rehman, A., Ullah, N., Khan, J., Qamar, N., ... & Khan, Y. (2021). Evaluation of change in hematological parameters and epidemiological identification of dengue virus infection at district Peshawar, Khyber Pakhtunkhwa, Pakistan. *International Journal of Mosquito Research*, 8(1, Part A), 11-18.
21. Anwar, F. (2020). Serological and Epidemiological Evaluation of active HCV Infection in the Volunteer blood Donor at District Swat Khyber Pakhtunkhwa Pakistan. *Bull. Env. Pharmacol. Life Sci*, 9,08-15

22. Bashir, Z., Ahmad, S. U., Kiani, B. H., Jan, Z., Khan, N., Khan, U., ...& Mahmood, T. (2021). Immuno informatics approaches to explore B and T cell epitope-based vaccine designing for SARS-CoV-2 Virus. *Pakistan Journal of Pharmaceutical Sciences*, 34
23. Ahmad, S. U., Khan, M. S., Jan, Z., Khan, N., Ali, A., Rehman, N., ...&Zahir, F. (2021). Genome wide association study and phylogenetic analysis of novel SARS-COV-2 virus amongdifferentcountries. *Pakistan Journal of Pharmaceutical Sciences*, 34(4)