

Assessment of Knowledge and Practices of Health Care Workers towards Infection Safety Precautions and Associated Factors in Health Care Facilities in the Makkah AL-Mukarramah, Saudi Arabia2022

Hassan Muwafak Habis¹, Meshal Mossad Alsaahfi¹, Badra Abdullah Alotaibi², Fouad Mohammed Alshehri³, Walid Saud Al Saeidi⁴, Azza Anas Abodawood⁵, Ahmad Abdulaziz Alhazmi⁶, Sami Aboud Alotaibi⁷, Sharea Thawab Alqahtani⁸, Abdulsalam Hassan Hassan Abu Sulayman⁹, Mansour Ali Zughbi¹⁰, Khalid Othman Haris Alzahrani¹¹, Rahma Atieh Eisa Alzahrani¹², Torki Saud Sad Alharthi¹³, Mohammed Ahmed Sari Alsuhaymi¹⁴

¹Doctor, King Abdalaziz Hospital Makkah, Saudi Arabia

²Family medicine, Primary health care 7 alsharyea, Makkah, Saudi Arabia

³Health services and hospital management specialist Aljumom health sector Ministry of Health, Makkah, Saudi Arabia

⁴Nursing Technician, Aljumom health sector Ministry of Health, Makkah, Saudi Arabia.

⁵Dentist, King Abdullah medical H jeddah - Alwafa polyclinic, Saudi Arabia.

⁶Epidemiologist, Department of Environmental and Occupational Health, Saudi Arabia.

⁷specialist Nursing, Hospital alnoor specialist, Makkah, Saudi Arabia.

⁸Pharmacist, King Abdulaziz Hospital, Makkah, Saudi Arabia.

⁹Pharmacy Technicians, King Abdulaziz Hospital, Makkah, Saudi Arabia.

¹⁰General Physician, King Abdulaziz Hospital, Makkah, Saudi Arabia.

¹¹Medical secretary technician, Maternity and children Hospital, Makkah, Saudi Arabia.

¹²Nurse, Health Center Al Kor Shadad, Makkah, Saudi Arabia.

¹³Epidemiology Technician, Primary health care center, Makkah, Saudi Arabia.

¹⁴Specialist Epidemiology, King Abdalaziz Hospital Makkah, Saudi Arabia.

Abstract

Background

The prevention of healthcare infection safety precautions and associated factors in health care facilities the provision of safe, high quality healthcare. Infections acquired in healthcare facilities are a major public health concern, contributing to increased morbidity, mortality, and cost in both developed and developing countries. Although most of these infections can be avoided by infection safety precautions with relatively inexpensive infection and control measures in many developing countries, in Saudi Arabia healthcare facilities have no effective infection safety precautions . Additionally, there is limited information on healthcare worker infection safety precautions knowledge and practice in countries such as Saudi Arabia .

Aim of study: To Assessment of knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities in the Makkah AL-Mukarramah, Saudi Arabia2022.

Methods: This cross sectional study included (200) health care professionals in Makkah City at Saudi Arabia 2023. (doctors, nurses, lab workers) from primary healthcare (PHC) centers an self-administrated questionnaire was constructed by the researcher and was used for data collection. Divided in to 3 parts i.e., socio-demographic characteristics, knowledge questions about infection safety precautions and statements about practice of health care providers regarding infection safety precautions and associated factors in health care facilities

Results: shows the majority of participant (55.0%) have **average** of the Knowledge about infection safety precautions , followed by (24.0%) of participant high but weak were (21.0%) while Range(2-9) and Mean \pm SD(7.12 \pm 1.99), while a significant relation were P-value=0.001 χ^2 63.78

Conclusion: Inadequate infection safety precautions knowledge and unsafe practices were frequent among study participants, reflecting a potentially common problem at public healthcare facilities in Saudi Arabia. Healthcare workers have better knowledge and safer practices if they had received infection safety precautions training and had infection safety precautions guidelines in their workplace. Interventions should be designed to consider these identified factors

Keywords: knowledge, practices, infection, safety, (HCWs) primary, health care, Makkah.

Introduction

Background

Workplace health and safety is a critical element in every organization particularly in the health care facilities [1]. Healthcare professionals (HCPs) may act as a mechanical vector for the transmission of infections from patient to patient [2, 3]. Accidental exposure to body fluids can cause blood borne infections particularly hepatitis B virus (HBV), human immunodeficiency virus (HIV), and hepatitis C virus (HCV) [4]. Exposures may happen through needle stick or sharp injuries contaminated with infected body fluids or through contact with splashes [5]. The risk of Serco on version after percutaneous exposure to infected blood is approximately 0.1–0.3% for HIV, 2% for HCV, and 6–60% for HBV [6]. Infection safety precautions help to protect healthcare workers, patients, and visitors from health hazards [6–7]. It includes the use of personal protective equipment (PPE) including gloves, gowns, eye goggles, aprons, masks [8], and through the provision of professional immunization programs [9]. With inadequate infection safety precautions , the risk of acquiring infections through exposure to body fluids is substantial [10]

According to the Centers for Disease Control and Prevention (CDC), HAIs defined as infections localized or systemic condition resulting from adverse reaction to the presence of infectious agent or its toxins acquired from health care settings that was not incubating or symptomatic at the time of admission to the healthcare facility [11]. These infections are a major public health concern and a threat to patient safety, contributing to increased morbidity, mortality, and cost [12]. Based on the available evidence, the overall impact of HAIs implies prolonged in the PHC stay, long-term disability, increased resistance of microorganisms to antimicrobials, high costs for patients and their family, and unnecessary deaths [13]. In addition, it places a significant massive additional economic burden on the health care system [14]. Many HCAIs are caused by micro- organisms present on the patient's body (resident flora) or from transient sources such as healthcare workers' hands,

contaminated equipment, or the environment.[15] The spread of these organisms usually results from breaches of core components of infection safety precautions measures, such as hand hygiene, disinfection and sterilization of instruments, and use of Personal infection safety precautions, safe disposal of wastes, sharps, and handling soiled linen as well as personal health and safety education, immunization programs, and post-exposure prophylaxis.[16] The use of effective and infection safety precautions measures possibly reduces the burden of HCAs by at least 30%.[17]

Despite the availability of these low-cost IPC strategies, compliance with standard IPC practices remains very low, especially in low- and middle-income countries.[18]

Globally, HCAs affect hundreds of millions of people each year.[19] Over 90% of these infections occur in resource- limited countries, mostly in Africa, where infections are more prevalent and adherence to infection safety precautions weak.[20] In addition, 15% of healthcare wastes are considered hazardous,16 which subsequently results in adverse healthcare outcomes such as prolonged PHC stays, long-term disability, increased antimicrobial resistance, massive additional financial burden on healthcare systems, high costs to patients and their families, and unnecessary deaths.[21]

Literature review

Faith , et al.(2019) study in Nigeria about good and fair knowledge among participants was reported as 50% and 44% respectively.[22] In Ethiopia, Yakob et al. showed that all participants had acceptable knowledge about contaminated needles and sharp materials that transmit disease causative agents, while 70.4% knew that gloves and gowns were required for any contact with patients.[23] In Brazil, Oliveria et al. identified a gap between knowledge of standard precautions and the practical applications among physicians.[24] In Ethiopia (2019) Beyamo et al assessed the compliance of health care workers with standard precaution practices and identified its determinants in public health institutions. The study included 250 HCWs. Nearly two-thirds (65%) of them had complied with standard precaution practices. Factors significantly associated with compliance to standard precaution practices were experience of ≤ 5 years, training on standard precaution, having good hand hygiene and availability of (personal protective equipment's)[25]

Online searching for studies exploring the knowledge and practices of health care workers infection safety precautions among primary healthcare workers yielded relatively few studies as most studies conducted in this field were among healthcare workers in hospitals and future health care workers. In addition, relatively limited studies were carried out in Saudi Arabia. [26]

In Al-Kharj, Alotaibi et al assessed the knowledge of as well as compliance of health care students with standard precautions. Results revealed that among surveyed 353 students, 70% had previously attended an infection control course. The knowledge and compliance with SPs levels were high. The commonest source of information self-learning while the current curriculum was the least reported one. Female students were more knowledgeable and compliant with SPs compared to males. Student`s specialty and academic level were significantly associated with knowledge and compliance regarding SPs. [26]

In Al-Qassim (2018), Al Ra`awji et al evaluated in a multicenter cross-sectional study among 354 HCWs the knowledge, attitudes, and practices regarding guidelines of hand hygiene. The

average knowledge score was 63%. Health-care workers aged over 30 years had higher scores than those younger than 30 years. Those at tertiary care hospitals had higher scores than those at secondary hospitals. Almost all had positive attitudes toward hand hygiene as well as adhering to the guidelines regularly. This study concentrated on only hand hygiene as a component of standard precautions. [27]

In Makkah, Alkot et al (2016) assessed the knowledge, attitude, and practice of health care workers toward Middle East respiratory syndrome coronavirus (MERS-CoV) among HCWs in primary health-care centers after an interventional education program. The level of satisfactory knowledge, positive attitude, and good practice of studied HCWs were significantly improved after exposure to the program, as it increased from 43.3%, 45%, and 57.4% before intervention to 67.9%, 63.8%, and 64.8% after intervention, respectively ($P < 0.001$). Older age, previous training, and experience were positively correlated with higher scores of knowledge.[28]

In Hofuf (2013), Amin et al (2013) evaluated in a cross-sectional study the knowledge of clinical years medical students about standard precautions of infection control' and explored their attitudes toward the current curriculum and training in providing them with effective knowledge and needed skills with this regard. The study included 251 students. Approximately one-quarter of them (26.7%) scored ≥ 24 (out of 41points) which was considered as an acceptable level. The least knowledge score were in the areas of sharp injuries, personal protective equipment and health care of the providers. The main sources of information were self-learning, and informal bed side practices. The majority of the participants believed that the current teaching and training regarding standard precautions are insufficient in providing them with the required knowledge and skills. These studies targeted future healthcare workers. [29]

Rationale

The Previous study found that almost half of the health care workers had poor towards infection safety precautions. The finding underlines the importance of good towards infection safety precautions knowledge and the accessibility of towards infection safety precautions guidelines to improve towards infection safety precautions practices among healthcare workers staff. The findings of this study also assessment of knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities in the Makkah AL-Mukarramah, Saudi Arabia. Effective knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities as well as practicing them properly is very critical in controlling the transmission infections among HCWs, The difficult challenge faced by the Saudi Ministry of Health is the healthcare services. Services that are provided free of charge to all Saudi citizens, increasing awareness of health and disease..

Aim of the study

To assessment of knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities in the Makkah AL-Mukarramah, Saudi Arabia2022

General objective:

To assessment of knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities in the Makkah AL-Mukarramah, Saudi Arabia2022.

Materials and methods .

Study design:

This study is descriptive cross-sectional study

Study sitting:

The study has been carried out in the city of Makkah Al-Mokarramah Makkah PHC centers in in Makkah City at Saudi Arabia Region. There are 300 primary health care centers belonging to Ministry of health (MOH) distributed as North (20) and South (19)

Study population:

MOH PHC health care professionals (n=300) distributed as follows: 50 physicians, nurses and laboratory technicians, Dental assistant, Dentist Study duration: September to October 2022 .

Sample size:

Sample size was calculated using open Epi online sample size calculator at 95% confidence level with bound on error of 5% regarding standard infection control precautions max sample size required is 300 participants.

Sample technique:

Sample technique was two stage.

At first stage: simple random sampling method will be used to select primary health care centers. At second stage: all the doctors, nurses and laboratory technicians within the selected PHCCs enrolled in the study. There are total primary health care centers. Expected numbers of HWs per each center are 10. So, we need 20 centers to collect the sample size.

Inclusion criteria:

Primary health care workers (doctors, nurses, laboratory technicians) in PHC center male and female, Saudi and non-Saudi, all ages, those who agreed to participate in the research.

Exclusion criteria:

Pharmacists, dentists, dental assistant. Those who have Vacation, disabled and absent during the data collection period .

Data collection tool and technique:

Data were collected by self-administrated questionnaire.

First part of the questionnaire includes questions about Demographic data of the physicians (gender, age, nationality, job title)

Second part about knowledge, and practice of towards infection safety precautions which including hand will be assessed covering hand hygiene obtained from WHO injection safety, and protective equipment utilization with barriers of adherence to towards infection safety precautions .

Data analysis:

Data were entered and analyzed using Statistical Package for Social Sciences (SPSS) software, version 24. Descriptive analysis was carried out as the mean and standard deviation

(SD) were calculated for quantitative variables, frequency and proportion were calculated for categorical variables.

For comparisons, chi-square and t-test was used for categorical and quantitative variables respectively. p –value ≤ 0.05 was considered significant for all inferential analysis.

Ethical approval:

- The ethical approval was taken from the Regional Research Ethics committee. A permission letter was obtained from the regional director of the city of Makkah Al-Mokarramah Makkah MOH before starting the data collection.
- A written Informed consent was obtained from each participant from commencing the data collection.
- The researcher preserved the confidentiality of the participants at all steps of the study for the data collection, analysis and result.

Budget: Self-funded.

Table 1 Distribution of Socio-demographic characteristics of Personal characteristics of the participants (n=300)

| | N | % |
|-----------------------|-----|----|
| Age | | |
| <20 years | 57 | 19 |
| 30-39 years | 111 | 37 |
| 40 -49 years | 87 | 29 |
| < 50 | 45 | 15 |
| Gender | | |
| Female | 132 | 44 |
| Male | 168 | 56 |
| Nationality | | |
| Non-Saudi | 33 | 11 |
| Saudi | 267 | 89 |
| Marital status | | |
| Single | 111 | 37 |
| Married | 132 | 44 |
| Divorced | 36 | 12 |
| Widowed | 21 | 7 |
| Profession | | |
| Physician | 63 | 21 |
| Dentist | 39 | 13 |
| Nurse | 123 | 41 |
| Lab technician | 33 | 11 |
| Dental assistant | 42 | 14 |
| Qualification | | |
| PhD/MD/equivalent | 51 | 17 |
| Master | 75 | 25 |

| | | |
|--|-----|-----------|
| Bachelor | 138 | 46 |
| Diploma | 36 | 12 |
| Experience in PHC | | |
| <5 years | 48 | 16 |
| 5-10 years | 135 | 45 |
| >10 years | 117 | 39 |
| Ever taken training on safety precautions | | |
| Yes | 237 | 79 |
| No | 63 | 21 |
| Presence of safety precautions committee | | |
| Yes | 243 | 81 |
| No | 57 | 19 |
| Availability of safety precautions guidelines in the working department | | |
| Yes | 207 | 69 |
| No | 93 | 31 |

Table 1 shows there were 300 participants, and the majority age was(37.0%) in (30-39)years, while the age 40-49 years were(29.0%) followed by <20 years were (19.%) the majority of them were males (56.0%) while female(44.0%), regarding the Nationality most of participants Saudi were(89.0%) but the non- Saudi were (11.0%), regarding marital status the majority of participant married were(44.0%)followed by single were(37.0%) followed by divorced were(12.0%), regarding profession the majority of participant are nurse were(41.0%) followed by Physician were(21.0%) followed by dentist were(14.0%), regarding qualification the majority of participant are bachelor were (46.0%) followed by master were(25.0%) followed by PhD/MD/equivalent were(17.0%), regarding experience in PHC the majority of participant are 5-10 years were (45.0%)followed by >10 years were(39.0%) followed by <5 years were(16.0%), regarding the ever taken training on safety precautions most of participants answer Yes were(79.0%) followed by No were(21.0%), regarding the presence of safety precautions committee the majority of participant answer Yes were (81.0%) followed by No were(19.0%), regarding the Availability of safety precautions guidelines in the working department the majority of participant answer Yes were (69.0%) followed by No were(31.0%) .

Table 2: Distribution of the Healthcare workers knowledge regarding infection safety precautions .

| Knowledge items | Yes | | No | | Chi-Square | |
|--|-----|----|-----|----|----------------|---------|
| | N | % | N | % | X ² | P-value |
| I have heard about infection safety precautions | 282 | 94 | 18 | 6 | 232.320 | <0.001* |
| Gloves cannot provide complete safety protection against transmission of infections | 216 | 72 | 84 | 28 | 58.080 | <0.001* |
| Washing hands with soap or use of an alcohol based antiseptic decreases the risk of transmission of healthcare acquired infections | 87 | 29 | 213 | 71 | 52.920 | <0.001* |
| Use of an alcohol based antiseptic for hand hygiene is as effective as soap and water if hands are not visibly dirty | 207 | 69 | 93 | 31 | 43.320 | <0.001* |
| Gloves should be worn if blood or body fluid exposure is anticipated | 264 | 88 | 36 | 12 | 173.280 | <0.001* |
| Hand washing is necessary before procedures are performed | 279 | 93 | 21 | 7 | 221.880 | <0.001* |
| Tuberculosis (TB) is carried in airborne particles that are generated from patients with active pulmonary tuberculosis | 240 | 80 | 60 | 20 | 108.000 | <0.001* |
| There is no need to change gloves between patients as long as there is no visible | 291 | 97 | 9 | 3 | 265.080 | <0.001* |

| | | | | | | |
|---|-----|----|----|----|---------|---------|
| contamination | | | | | | |
| Do you know how to prepare 0.5% chlorine solution | 273 | 91 | 27 | 9 | 201.720 | <0.001* |
| Safety box should be closed/sealed when three quarters filled | 264 | 88 | 36 | 12 | 173.280 | <0.001* |

Table 2 shows the distribution of the Healthcare workers knowledge regarding I have heard about infection safety precautions, the majority of participant answer Yes were (94.0%) while answer No were (6.0%), while is a significant relation were P-value=0.001 χ^2 232.320, regarding Gloves cannot provide complete safety protection against transmission of infections, the majority of participant answer Yes were (72.0%) while answer No were (28.0%), while is a significant relation were P-value=0.001 χ^2 58.080, regarding washing hands with soap or use of an alcohol based antiseptic decreases the risk of transmission of healthcare acquired infections, the majority of participant answer No were (71.0%) while answer Yes were (29.0%), while is a significant relation were P-value=0.001 χ^2 52.920, regarding Use of an alcohol based antiseptic for hand hygiene is as effective as soap and water if hands are not visibly dirty the majority of participant answer Yes were (69.0%) while answer No were (31.0%), while is a significant relation were P-value=0.001 χ^2 43.320, regarding gloves should be worn if blood or body fluid exposure is anticipated, the majority of participant answer Yes were (88.0%) while answer No were (12.0%), while is a significant relation were P-value=0.001 χ^2 173.280, regarding hand washing is necessary before procedures are performed the majority of participant answer Yes were (93.0%) while answer No were (7.0%), while is a significant relation were P-value=0.001 χ^2 108.000, regarding Tuberculosis (TB) is carried in airborne particles that are generated from patients with active pulmonary tuberculosis the majority of participant answer Yes were (80.0%) while answer No were (2.0%), while is a significant relation were P-value=0.001 χ^2 108.000, regarding There is no need to change gloves between patients as long as there is no visible contamination the majority of participant answer Yes were (97.0%) while answer No were (3.0%), while is a significant relation were P-value=0.001 χ^2 265.080, regarding Do you know how to prepare 0.5% chlorine solution the majority of participant answer Yes were (91.0%) while answer No were (9.0%), while is a significant relation were P-value=0.001 χ^2 201.720, regarding Safety box should be closed/sealed when three quarters filled the majority of participant answer Yes were (88.0%) while answer No were (12.0%), while is a significant relation were P-value=0.001 χ^2 173.280,

Table 3: Distribution of the Healthcare workers Practice towards infection safety precautions .

| Practice items | Yes | | No | | Chi-Square | |
|--|-----|----|-----|----|----------------|---------|
| | N | % | N | % | X ² | P-value |
| Do you apply antiseptic hand rub to clean hands? | 225 | 75 | 75 | 25 | 75.000 | <0.001* |
| Did you practice high-level disinfection where sterilization is not applicable? | 216 | 72 | 84 | 28 | 58.080 | <0.001* |
| Do you use all Personal Protective Equipment's (PPE) to prevent the risk of acquiring and /or transmitting infection? | 198 | 66 | 102 | 34 | 30.720 | <0.001* |
| Did you mix dry and liquid Healthcare wastes? | 201 | 67 | 99 | 33 | 34.680 | <0.001* |
| Do you incinerate or bury used sharp materials? | 207 | 69 | 93 | 31 | 43.320 | <0.001* |
| You change disinfectant chlorine solutions? | 225 | 75 | 75 | 25 | 75.000 | <0.001* |
| Do you know how long it takes to soak reusable medical instruments in chlorine Solution? | 213 | 71 | 87 | 29 | 52.920 | <0.001* |
| Do you use gloves more than once (both hands)? | 207 | 69 | 93 | 31 | 43.320 | <0.001* |
| Do you wear the necessary personal protective equipment (PPE) such as gloves, apron, goggles and mask, if splashes and spills of any body fluids are likely? | 243 | 81 | 57 | 19 | 115.320 | <0.001* |
| Where do you usually put sharp disposal boxes? | 237 | 79 | 63 | 21 | 100.920 | <0.001* |

Table 3 shows the distribution of the healthcare workers Practice towards infection safety precautions regarding you apply antiseptic hand rub to clean hands the majority of participant answer Yes were (75.0%) while answer No were (25.0%), while is a significant relation were P-value=0.001 X² 58.080, regarding you practice high-level disinfection where sterilization is not applicable the majority of participant answer Yes were (72.0%) while answer No were (28.0%), while is a significant relation were P-value=0.001 X² 58.080, regarding you use all Personal Protective Equipment's to prevent the risk of acquiring and /or transmitting infection the majority of participant answer Yes were (66.0%) while answer No were (34.0%), while is a significant relation were P-value=0.001 X² 30.720, regarding you mix dry and liquid Healthcare wastes the majority of participant answer Yes were (67.0%) while answer No were (33.0%), while is a significant relation were P-value=0.001 X²

34.680, regarding you incinerate or bury used sharp materials, the majority of participant answer Yes were (69.0%) while answer No were (31.0%), while is a significant relation were P-value=0.001 χ^2 43.320, regarding you change disinfectant chlorine solutions the majority of participant answer Yes were (75.0%) while answer No were (25.0%), while is a significant relation were P-value=0.001 χ^2 52.920, regarding you know how long it takes to soak reusable medical instruments in chlorine Solution the majority of participant answer Yes were (71.0%) while answer No were (29.0%), while is a significant relation were P-value=0.001 χ^2 52.920, regarding you use gloves more than once (both hands) the majority of participant answer Yes were (69.0%) while answer No were (31.0%), while is a significant relation were P-value=0.001 χ^2 43.320, regarding you wear the necessary personal protective equipment (PPE) such as gloves, apron, goggles and mask, if splashes and spills of any body fluids are likely the majority of participant answer Yes were (81.0%) while answer No were (21.0%), while is a significant relation were P-value=0.001 χ^2 100.920, regarding Where do you usually put sharp disposal boxes the majority of participant answer Yes were (79.0%) while answer No were (21.0%), while is a significant relation were P-value=0.001 χ^2 100.920,

Table 4: Distribution the knowledge of Healthcare workers towards infection safety precautions .

| | Knowledge | | Score | |
|----------------------------|-----------|-----|-------|-----------|
| | N | % | Range | Mean±SD |
| Weak | 63 | 21 | 2-9. | 7.12±1.99 |
| Average | 165 | 55 | | |
| High | 72 | 24 | | |
| Total | 300 | 100 | | |
| χ^2 | 63.78 | | | |
| P-value | <0.001* | | | |

This table 4 shows the majority of participant (55.0%) have **average** of the Knowledge about infection safety precautions , followed by (24.0%) of participant high but weak were (21.0%) while Range(2-9) and Mean ±SD(7.12 ±1.99), while a significant relation were P-value=0.001 χ^2 63.78 .

Figure (1): Distribution the knowledge of Healthcare workers towards infection safety precautions




Table 5: Distribution the Practices of Healthcare workers towards infection safety precautions

| | Practices | | Score | |
|----------------------|-----------|-----|-------|-----------|
| | N | % | Range | Mean±SD |
| Weak | 117 | 39 | 1-9. | 5.64±1.75 |
| Average | 132 | 44 | | |
| High | 51 | 17 | | |
| Total | 300 | 100 | | |
| X² | 37.14 | | | |
| P-value | <0.001* | | | |

regarding the practice the majority of participant (44. 0%) have average of the practice about infection safety precautions, followed by (39.0%) of participant weak while high were(17.0)while Range(1-9) and Mean ±SD(5.64 ±1.75) While a significant relation were P-value=0.001 X² 37.14

Figure (2): Distribution the Practices of Healthcare workers towards infection safety precautions




Table 6: Distribution of the Correlations between Knowledge and Practices of Healthcare workers towards infection safety precautions

| Correlations | | |
|------------------|----------------|------------------|
| | | Knowledge |
| Practices | r | 0.532 |
| | P-value | <0.001* |
| | N | 300 |

This table 6 shows the heave correlations between Knowledge and Practices while r were 0.532 and a significant relation were P-value=0.001 and N(300)

Figure (3): Distribution of the Correlations between Knowledge and Practices of Healthcare workers towards infection safety precautions Discussion .



Reducing the risk of HAIs and using infection safety precautions principles are in the control of healthcare workers; therefore, healthcare workers must have correct, up-to-date and appropriate scientific information and practice accordingly [30]. Without knowledge infection safety precautions and patient safety practices both healthcare workers and patients are at risk of acquiring serious infections such as HIV, HBV, HCV, and Methicillin Resistant *Staphylococcus aureus* (MRSA) infection as well as other bacterial and viral infections [27]. Recent studies also suggest that proper and consistent application of infection safety precautions and infection safety practices can lead to up to a 70% reduction in certain HAIs [31]. This study was conducted to assessment of knowledge and practices of health care workers towards infection safety precautions and associated factors in health care facilities in the Makkah AL-Mukarramah at the primary health care level. The of knowledge and practices of health care workers towards infection safety precautions is an essential step in starting and implementing a successful infection safety precautions in any healthcare facility.[32] Worldwide, many studies have shown that healthcare workers expressed variable levels of knowledge regarding infection safety precautions, with relatively limited studies have been carried out in the Kingdom of Saudi Arabia.[33] Therefore, the present study was conducted the present study. One of the most important characteristics of Makkah is its location, which is characterized by proximity to Makkah. In our study showed were 300 participants, and the majority age was(37.0%) in (30-39)years, males (56.0%) most of participants Saudi were(89.0%), profession the majority of participant are nurse were(41.0%). regarding experience in PHC the majority of participant are 5-10 years were (45.0%), regarding the presence of safety precautions committee the majority of participant answer Yes were (81.0%) (see Table 1) Also showed that most of participants had high knowledge regarding infection safety precautions. In this study, the proportion of healthcare workers who were knowledgeable about infection safety precautions majority are average was found to be 55.0%. similar studies in Ethiopia [30]]This finding indicated that a large percentage of respondents (46.3%) in the healthcare infection safety precautions and associated factors in health care facilities inadequate knowledge about infection prevention, a finding in line with similar studies in Africa [31]. (See table 4,5)

On the other hand, the proportion of knowledgeable participants is lower than studies in facilities in Bahir Dar city and Addis Ababa which reported 69% and 84.2% of healthcare workers had good infection safety precautions knowledge [33]

This discrepancy may be due to difference in study setting and study variables since the former study focused only the two components of infection prevention (hand hygiene and tuberculosis infection control) and includes only two university hospitals in Addis Ababa the later includes private healthcare facilities. Similarly, the result is inconsistent with that. [34].

Conclusion

The present study revealed that a significant proportion of healthcare workers were knowledgeable about infection safety precautions. The overall level of safe infection safety precautions practice among healthcare workers is considered to be weak. The current study also detected that there was a high prevalence of occupational needle stick injury and blood and body fluid splashes among healthcare workers. Factors such as the presence of infection prevention guidelines in the work place and training were independent predictors of infection

safety precautions practice and better knowledge. Providing on job continuous educational training on infection safety precautions is essential as well as ensuring the availability of infection safety precautions guidelines in working department should be effective and important interventions to improve healthcare workers infection safety precautions and knowledge. In the future researchers should consider stronger observational study designs to validate the self-reported practice of healthcare workers and to determine actual practices, as well as the actual prevalence of HAIs as result of poor infection safety precautions practice

References

- [1] Somani, R., Muntaner, C., Hillan, E., Velonis, A. J., & Smith, P. (2021). A systematic review: effectiveness of interventions to de-escalate workplace violence against nurses in healthcare settings. *Safety and health at work*, *12*(3), 289-295.
- [2] Latha, T., Bhat, A. K., Hande, H. M., Mukhopadhyay, C., Devi, E. S., Nayak, B. S., & George, A. (2022). Effectiveness of Extended Infection Control Measures on Methicillin-Resistant Staphylococcus aureus Infection Among Orthopaedic Patients. *Indian Journal of Orthopaedics*, *56*(10), 1804-1812.
- [3] Hayden, J. (2022). *Introduction to health behavior theory*. Jones & Bartlett Learning.
- [4] Alhumaid, S., Al Mutair, A., Al Alawi, Z., Alsuliman, M., Ahmed, G. Y., Rabaan, A. A., ... & Al-Omari, A. (2021). Knowledge of infection prevention and control among healthcare workers and factors influencing compliance: a systematic review. *Antimicrobial Resistance & Infection Control*, *10*(1), 1-32.
- [5] Roien, R., Mousavi, S. H., Ozaki, A., Baqeri, S. A., Hosseini, S. M. R., Ahmad, S., & Shrestha, S. (2021). Assessment of knowledge, attitude, and practice of health-care workers towards hepatitis B virus prevention in Kabul, Afghanistan. *Journal of Multidisciplinary Healthcare*, 3177-3186.
- [6] Sahiledengle, B., Tekalegn, Y., & Woldeyohannes, D. (2021). The critical role of infection prevention overlooked in Ethiopia, only one-half of health-care workers had safe practice: A systematic review and meta-analysis. *PloS one*, *16*(1), e0245469.
- [7] AlJohani, A., Karuppiah, K., Al Mutairi, A., & Al Mutair, A. (2021). Narrative review of infection control knowledge and attitude among healthcare workers. *Journal of epidemiology and global health*, *11*(1), 20.
- [8] Sabetian, G., Moghadami, M., Hashemizadeh Fard Haghghi, L., Shahriarirad, R., Fallahi, M. J., Asmarian, N., & Moeini, Y. S. (2021). COVID-19 infection among healthcare workers: a cross-sectional study in southwest Iran. *Virology journal*, *18*, 1-8.
- [9] Tsegaye, D., Shuremu, M., Oljira, D., Dubale, S., Befekadu, G., & Bidira, K. (2021). COVID-19 related knowledge and preventive practices early in the outbreak among health care workers in selected public health facilities of Illu aba Bor and Buno Bedelle zones, Southwest Ethiopia. *BMC Infectious Diseases*, *21*(1), 1-11.
- [10] Ilesanmi, O. S., Afolabi, A. A., Akande, A., Raji, T., & Mohammed, A. (2021). Infection prevention and control during COVID-19 pandemic: realities from health care workers in a north central state in Nigeria. *Epidemiology & Infection*, *149*, e15.
- [11] Abuduxike, G., Vaizoglu, S. A., Asut, O., & Cali, S. (2021). An assessment of the knowledge, attitude, and practice toward standard precautions among health workers from a hospital in northern cyprus. *Safety and health at work*, *12*(1), 66-73.

- [12] Lai, T. H., Tang, E. W., Chau, S. K., Fung, K. S., & Li, K. K. (2020). Stepping up infection control measures in ophthalmology during the novel coronavirus outbreak: an experience from Hong Kong. *Graefe's Archive for Clinical and Experimental Ophthalmology*, 258(5), 1049-1055.
- [13] Ochie, C. N., Aniwada, E. C., Uchegbu, E. K., Asogwa, T. C., & Onwasoigwe, C. N. (2022). Infection prevention and control: knowledge, determinants and compliance among primary healthcare workers in enugu metropolis, south-east nigeria. *Infection Prevention in Practice*, 4(2), 100214.
- [14] Chigurupati, R., Panchal, N., Henry, A. M., Batal, H., Sethi, A., D'innocenzo, R., ... & Roser, S. M. (2020). Considerations for oral and maxillofacial surgeons in COVID-19 era: can we sustain the solutions to keep our patients and healthcare personnel safe?. *Journal of Oral and Maxillofacial Surgery*, 78(8), 1241-1256.
- [15] Curry, S. R., & Salgado, C. D. (2021). When hospitals harm: multimodal entry of SARS-CoV-2 into inpatient healthcare. *Clinical Infectious Diseases*, 72(4), 694-696..
- [16] Asdaq, S. M. B., Alshrari, A. S., Imran, M., Sreeharsha, N., & Sultana, R. (2021). Knowledge, attitude and practices of healthcare professionals of Riyadh, Saudi Arabia towards covid-19: A cross-sectional study. *Saudi Journal of Biological Sciences*, 28(9), 5275-5282.
- [17] Al-Ahmari, A. M., AlKhalidi, Y. M., & Al-Asmari, B. A. (2021). Knowledge, attitude and practice about infection control among primary care professionals in Abha City, Kingdom of Saudi Arabia. *Journal of Family Medicine and Primary Care*, 10(2), 662
- [18] Haines, A., de Barros, E. F., Berlin, A., Heymann, D. L., & Harris, M. J. (2020). National UK programme of community health workers for COVID-19 response. *The Lancet*, 395(10231), 1173-1175.
- [19] Kung, S., Doppen, M., Black, M., Hills, T., & Kearns, N. (2021). Reduced mortality in New Zealand during the COVID-19 pandemic. *The Lancet*, 397(10268), 25.
- [20] Lancet, T. (2020). COVID-19: learning from experience. *Lancet (London, England)*, 395(10229), 1011.
- [21] Albarrak, A. I., Mohammed, R., Al Elayan, A., Al Fawaz, F., Al Masry, M., Al Shammari, M., & Miaygil, S. B. (2021). Middle East Respiratory Syndrome (MERS): Comparing the knowledge, attitude and practices of different health care workers. *Journal of infection and public health*, 14(1), 89-96.
- [22] Faith, I., Harrison, E., Aigbiremolen Alphonsus, O., Ekundare, F. O., Rowland-Udoh Eloho, A., Ogeyemhe Charles, O., & Okudo Ifeanyi, O. A. B. (2019). Knowledge, attitude and infection prevention and control practices regarding Lassa fever among healthcare workers in Edo State, Nigeria. *Hospital*, 2, 2.
- [23] Faria, L. B. G. D., Santos, C. T. B. D., Faustino, A. M., Oliveira, L. M. D. A. C., & Cruz, K. C. T. D. (2019). Knowledge and adherence of the nurse to standard precautions in critical units. *Texto & Contexto-Enfermagem*, 28.
- [24] Oliveira, A. C., Marziale, M. H. P., Paiva, M. H. R. S., & Lopes, A. C. S. (2009). Knowledge and attitude regarding standard precautions in a Brazilian public emergency service: a cross-sectional study. *Revista da Escola de Enfermagem da USP*, 43, 313-319.

- [25] Beyamo, A., Dodicho, T., & Facha, W. (2019). Compliance with standard precaution practices and associated factors among health care workers in Dawuro Zone, South West Ethiopia, cross sectional study. *BMC health services research*, 19(1), 1-6.
- [26] Alotaibi, M. M., Almasari, S. M., Alkadam, A. N., Alanazi, Y. A., & Al Gahtani, K. A. (2017). Knowledge and compliance with standard isolation precautions among healthcare students in Al-Kharj Governorate, Saudi Arabia. *J Health Spec*, 5(3), 162-70.
- [27] Al Ra'awji, B. A., Almogbel, E. S., Alharbi, L. A., Alotaibi, A. K., Al-Qazlan, F. A., & Saquib, J. (2018). Knowledge, attitudes, and practices of health-care workers regarding hand hygiene guidelines in Al-Qassim, Saudi Arabia: A multicenter study. *International journal of health sciences*, 12(2), 3.
- [28] Alkot, M., Albouq, M. A., Shakuri, M. A., & Subahi, M. S. (2016). Knowledge, attitude, and practice toward MERS-CoV among primary health-care workers in Makkah Al-Mukarramah: an intervention study. *Int J Med Sci Public Health*, 5(5), 952-60.
- [29] Amin, T. T., Al Noaim, K. I., Saad, M. A. B., Al Malhm, T. A., Al Mulhim, A. A., & Al Awas, M. A. (2013). Standard precautions and infection control, medical students' knowledge and behavior at a Saudi university: the need for change. *Global journal of health science*, 5(4), 114.
- [30] Dalton, K. R., Rock, C., Carroll, K. C., & Davis, M. F. (2020). One Health in hospitals: how understanding the dynamics of people, animals, and the hospital built-environment can be used to better inform interventions for antimicrobial-resistant gram-positive infections. *Antimicrobial Resistance & Infection Control*, 9, 1-17.
- [31] Hosseinalhashemi, M., Kermani, F. S., Palenik, C. J., Poursaghari, H., & Askarian, M. (2015). Knowledge, attitudes, and practices of health care personnel concerning hand hygiene in Shiraz University of Medical Sciences hospitals, 2013-2014. *American journal of infection control*, 43(9), 1009-1011.
- [32] Tan, C., Kallon, I. I., Colvin, C. J., & Grant, A. D. (2020). Barriers and facilitators of tuberculosis infection prevention and control in low-and middle-income countries from the perspective of healthcare workers: A systematic review. *PloS one*, 15(10), e0241039
- [33] ALjohani, H. S., & Sulaiman, A. A. (2021). Assessment of Health Care Workers' Knowledge and Practice Toward Infection Standard Precautions in Primary Health Care setting, Buraidah, Saudi Arabia. *Middle East Journal of Family Medicine*, 7(10), 81.
- [34] ElBadry, S., Ghaleb, M. A., & Abou Zeid, N. A. (2019). Healthcare Personnel Opinion and their Implementation Obstacles Regarding the Standard Precautions in Hemodialysis Unit. *Evidence-Based Nursing Research*, 1(4), 13-13.