

¹Received 10 August 2023; Accepted 25 August 2023

Assessment of the Knowledge and Practices of Dengue Fever Symptoms Among Health Care Workers at the Primary Healthcare Level in Makkah City at Saudi Arabia 2023.

Emad Abdulaziz Almoalem¹, Sultan Hymid Al-Hazmi², Turki Ammar Omair Alhazmi³, Ibraheem Ahmed Alzahrani², Wisal Mohamad Noor Sinbawa⁴, Ayman Salem Ajyeeb⁵, Marwan Solimon Mouawad⁶, Hesham Abdullah Ganim⁷, Mohammed Meyassr Melybari⁸, Abdullah Fozy Almegren⁹, Hassan Fouad Labban⁹, Nahla Nasir Aljeaid¹⁰, Mashaal Musaad Mousa Albarakati¹¹, Jameelah Ghazi Allihaibi¹², Malak Yousef Ali¹²

¹Community medicine specialist, Department of Infectious Diseases Control in Makkah Health Cluster, Saudi Arabia

²Epidemiology Monitor, Department of Infectious Diseases Control in Makkah Health Cluster, Saudi Arabia

³Nursing Technician, Department of Infectious Diseases Control in Makkah Health Cluster, Saudi Arabia ⁴Nursing Technician, Alwali health center, Makkah, Saudi Arabia.

⁵Nursing Technician, Makkah Healthcare Cluster, MOH, Saudi Arabia.

⁶Technician of infection, Al Asila health center, Saudi Arabia.

⁷Nursing Technician, Alzahir Healthcare Cluster Makkah, MOH, Saudi Arabia.

⁸General Practitioner, Infection Control Directorate at Makkah Healthcare Cluster, Saudi Arabia.

⁹Nursing Technician, Infection Control Directorate at Makkah Healthcare Cluster, Saudi Arabia.

¹⁰Specialist nurse, 937 Department in the executive Operation Department at Makkah healthcare cluster, Saudi Arabia.

¹¹Specialist Social service, Ibn Sina hospital, Saudi Arabia.

¹²Internal medicine specialist, Alnoor specialist hospital Makkah, MOH, Saudi Arabia.

Abstract:

Background:

Dengue fever (DF) is a viral disease transmitted by mosquitoes and is emerging as a serious global health problem. However, there is still a lack of knowledge regarding the disease, to investigate the knowledge and practices of the healthcare professionals (HCPs) including physicians and nurses regarding dengue transmission, diagnosis and clinical classification using the warning signs of World Health Organization (WHO) classification, Dengue fever is a major arbovirus-borne infectious disease in tropical and subtropical regions of the world. The disease is generally mild and self-limited, but some patients may develop a severe form of infection such as dengue hemorrhagic fever/dengue shock syndrome, which often leads to death. Currently, there is no vaccine available against dengue virus. The major determinants of dengue control are vector eradication, early case recognition, and adequate clinical management, Knowledge and practice (KP) among primary healthcare professionals (HCPs) regarding dengue diseases may pace alarm and improve the outcome of dengue control, Primary physicians and nurses serve as the first-line health care providers of dengue virus infection diagnosis, notification, and treatment..

Aim of the study :To assess the Knowledge and practices of dengue fever symptoms among health care workers at the primary healthcare level in Makkah City at Saudi Arabia 2023.

¹Received 10 August 2023; Accepted 25 August 2023

Methods: Across sectional descriptive study conducted among health care workers selected primary health-care centers in Makkah Al-Mokarramah city, during the September to October, 2023, the Sample size of medical practitioners. Our total participants were (200).

Results: shows majority of participant were (34.0%) were aged from 30-45 years of age, sex, more than half of participant (58.0%) were male, experience the majority of participant <2 y were (43.0%), specialty the majority of participant internal medicine were (39.0%), majority of participant married were (49.0%) while single were(34.0%) but widow were (10.0%), regarding reported a dengue case more than half of participant answer Yes were (75.0%).

Conclusion: the most of the HCPs did not take prior training on the dengue viral infection; also, have gaps in the knowledge regarding various topics in dengue fever. This paper recommends the gathering of efforts to establish the proper knowledge of dengue infection and the warning signs listed by the WHO, educational programs for health care workers on dengue fever management and prevention should be organized.

Keywords: Dengue, fever, knowledge, practices, symptoms, health care workers, primary, Makkah .

Introduction

Dengue virus disease (DVD) that include dengue fever, dengue hemorrhagic fever and dengue shock syndrome, is caused by dengue virus; one of the Flaviviridae and it is considered by the World Health Organization (WHO) as the most rapid growing mosquito-borne infection and dengue virus can be co-transmitted with other viruses such as Zika viruses and human confessions have been reported (1,2). DVD has affected least 129 countries of which over 100 tropical and subtropical countries are considered endemic or hyper-endemic with an estimated global annual incidence up to 400 million annual infections of which only around (3) million infections were officially reported to the WHO in 2022. Furthermore, DVD is considered, probably due to climate change and/or viral evolutionary mutations, as an evolving threat to the remaining countries(4,5)

Dengue is not considered endemic in Saudi Arabia ; however, several outbreaks have been reported in southern Saudi Arabia.(6) Saudi Arabia dengue outbreaks have a unique type of transmission: starting by import from abroad in early summer, spreading locally, and ending in the winter. This pattern repeats every year.(7) Most dengue cases occurred in Saudi Arabia , thus, dengue control has been an important issue for healthcare professionals (HCPs) in these areas.(8)

Dengue is a viral disease transmitted to humans by the bite of infected females of the main vector *Aedes aegypti* and to lesser extent *Aedes albopictus* mosquitoes (9). The World Health Organization has classified dengue into three categories according to disease severity; Dengue Fever (DF), Dengue Hemorrhagic Fever (DHF), and Dengue Shock Syndrome (DSS)24. Severe dengue (DHF and DSS) causes lethal complications that included severe hemorrhage, plasma leakage, organ impairment, fluid accumulation, or respiratory distress .(10,11)

Disease Control and Prevention, there are two species of mosquitoes that transmit dengue fever, but the primary vector of dengue is *Aedes aegypti*. (12) This species, whose food source is human blood, lives mainly inside buildings in dark areas such as closets and

¹Received 10 August 2023; Accepted 25 August 2023

bathrooms. However, it can also be found in outdoor areas with standing water such as construction sites and gardens. (13)

Severe dengue (previously known as dengue hemorrhagic fever) was first recognized in the 1950s during dengue epidemics in the Philippines and Thailand. (15) Today, severe dengue affects most Asian and Latin American countries and has become a leading cause of hospitalization and death among children in these regions. (16) Dengue fever has a wide range of presentations from mild to severe. On the mild side, it entails a low, self-limited fever, but severe cases can entail life-threatening hemorrhagic shock. The incubation period of the dengue fever virus in humans ranges from 3 to 14 days. (17)

The World Health Organization (WHO) recognized the need for general guidelines for dengue management and published the first and revised editions of the dengue guideline in 1975 and 1997.(18) The 1975/1997 WHO guidelines divided clinical dengue disease into dengue fever, dengue hemorrhagic fever, and dengue shock syndrome.(19) Criticisms of the 1975/1997 guidelines included a rigid definition of dengue hemorrhagic fever and the lack of data from the adult group outside of the southeastern Asia region.(20) The WHO released the newest (third) edition of the dengue guideline in 2009 in response to the criticism after conducting the multicenter dengue control study.(21) By introducing the concepts of 'warning signs' and 'severe dengue', the 2009 guidelines intended to provide a more clinical management tool than merely a classification for dengue control and management.(22,23)

Literature Review

In 2009, the WHO published a classification of dengue cases, which was attributed to several benefits shown in literature (24). One of those values was the considerable sensitivity in capturing the severe case (25). Moreover, a study proved that this classification was easy to apply with its user-friendliness to healthcare providers (26). Nevertheless, the WHO 2009 classification listed seven warning signs of dengue without providing a precise definition of these signs. These lead to the different application of these signs for the diagnosis and treatment of dengue patients between healthcare personnel. (27)

In 2009, the Saudi MOH reported a total of 3350 cases of DF in the Kingdom and estimated the case fatality rate to be 4.6 per thousand (28). The reemergence of DF in Saudi Arabia can be explained by the growing levels of urbanization, international trade and travel .(29)

in another study found some major gaps in knowledge regarding some important issues that need reconsideration were identified such as believing that A Egypt, the mosquito vector for dengue virus, typically bites after dark, prescription of aspirin or ibuprofen for confirmed dengue case, dengue infection by one serotype DEN (30) will give lifelong immunity against all serotypes, and reporting that the most sensitive and specific method of acute dengue infection diagnosis is isolation in cell culture using immunofluorescence. Similar gaps in the PHC physicians' knowledge have been reported in a recent study conducted in Jeddah. (31)

Nanyingi (2018) reported, poor knowledge on transmission of dengue was evident from the respondent's inability to recognize the feeding time of dengue mosquitoes. Similar findings have been documented by Huang et al. in Taiwan, where, only 14.4% of respondents correctly identified the feeding behavior of dengue mosquitoes (32). Having accurate knowledge on the behavior of dengue mosquitoes will help MPs to impart appropriate health

¹Received 10 August 2023; Accepted 25 August 2023

education (33), which remains one of the cornerstones of preventing dengue. Respondents had low knowledge of the signs that lead to dengue shock and thrombocytopenia. Such findings were also reported in Puerto Rico, where only 29.0% of the respondents correctly identified early signs of shock, and 48.0% identified severe abdominal pain and persistent vomiting as warning signs of severe dengue (34).

Sahu et al.,(2023) report that dengue vector change occurs on a large range of intricate temporal and spatial scales, where the change occurs on a daily scale to where the evolution of a potential repetition zone of the vector occurs on a yearly scale. In addition, modeling the daily changes of hotspots of the vector is conceivable on a sub-district scale but not on a sub-municipality scale, where long-range interactions cannot be modeled accurately (30). This could be due to a lack of training on the recognition of warning signs and case classification of dengue as per the updated WHO guidelines. Identification of warning signs of dengue and indications that lead to shock is critical for managing dengue (24).

Yusuf and Ibrahim reported that 56.0% of respondents lacked adequate training to manage dengue patients, including identifying warning signs, and recommended to close this gap (35). also, previous knowledge, attitude and practices (KAP) studies concerning control of dengue virus showed the lack of knowledge about clinical features or control measures as the most common problem. (22) This study found almost a third of PHC physicians had insufficient knowledge about important investigations of dengue as well as prevention measures toward DF. In southern Taiwan, (36)

Rationale:

Since the first case of DHF died in Jeddah in 1993, Saudi Arabia has reported three major epidemics: a dengue virus serotype (DEN)-2 epidemics in 1994 with 469 cases of dengue, 23 of DHF. The incidence of DF in Saudi Arabia by 2020 was increase the percentage among Saudi nationals and also among non-Saudi nationals. Since 1994, Makkah province became a dengue-endemic area with high rate of dengue infection during spring and early summer. Several dengue outbreaks have been reported after 2015 in the Makkah region. Millions of Muslims living in hyper endemic dengue areas come yearly to Makkah for Hajj, with the possibility of introduction of dengue virus. Same time gaps in the health care workers knowledge have been reported in a recent study conducted also primary physicians and nurses serve as the first-line health care providers of dengue virus infection diagnosis, notification, and treatment. Knowledge, attitude, and practice (KAP) among primary healthcare professionals (HCPs) regarding dengue diseases may pace alarm and improve the outcome of dengue control. health education can be useful for DF surveillance systems for public health officials. This method can provide an opportunity to specify the health burden of DF.

2.1 Aim of the study:

To assess the Knowledge and practices of dengue fever symptoms among health care workers at the primary healthcare level in Makkah City at Saudi Arabia 2023

Objectives:

To assess the Knowledge and practices of dengue fever symptoms among health care workers at the primary healthcare level in Makkah City at Saudi Arabia 2023

¹Received 10 August 2023; Accepted 25 August 2023

3. Methodology:

3.1 Study design:

This study is a prospective cross-sectional study design was used in carrying out of this study.

3.2 Study Area

The study will be carried out in the city of Makkah Al-Mokarramah Makkah is the holiest spot on Earth. It is the birthplace of the Prophet Mohammad and the principal place of the pilgrims to perform Umrah and Hajj. It is located in the western area in Kingdom of Saudi Arabia and called the Holy Capital. Contains a population around 2.578 million. This study was conducted in Makkah primary health-care centers at Saudi Arabia, and it reflects a diversified demographic profile with a considerable portion of the population comes from rural descent, while others come from an urban one. This difference translates into biological, socioeconomic and lifestyle differences in the Makkah population.

3.3 Study Population

The study has be conducted among **health care workers in the primary healthcare** regarding the **Knowledge and practices of dengue fever symptoms** in Makkah the sample was selected to include primary health-care medical practitioners who aged from <30years - More than 60 years and their total number was 200

3.2.1 Selection criteria:

3.2.2 .Inclusion criteria

- All nationalities
- aged from 30 to 60 year

3.3.2Exclusion criteria :

- No specific exclusion criteria.

3.4 The sample size

The sample size has been calculated by applying Raosoft sample size calculator based on (The margin of error: 5%, Confidence level: 95%, and the response distribution was considered to be 20%) accordingly to sample size from medical practitioners by the required sample size; (200). (male and female) and adding 10 more to decrease margin of error. After adding 5% oversampling, the minimum calculated sample has been 200. Computer generated simple random sampling technique was used to select the study participants. Data collection was done by the researcher during the September to October to, 2023.

Sampling technique:

Systematic random sampling technique is adopted. After that, by using random number generator, then simple random sampling technique was applied to select the medical practitioners. Also, convenience sampling technique will be utilized to select the participants in the study. By using systematic sampling random as dividing the total medical practitioners by the required sample size; (200).

Data collection tools of the study:

Tool was designed to collect the necessary data, and developed by the researchers after review of the literature.

Tool I: Health care workers knowledge and practices regarding dengue fever structured interview questionnaire:

It included five parts as follows:

¹Received 10 August 2023; Accepted 25 August 2023

Part one: Health care workers socio demographic characteristics:

This part consisted data about patient's age, sex, marital status, level of education, religion, income and sources of information.

Part two: Knowledge about signs and symptoms of dengue fever:

Include items that determine the health care workers knowledge about clinical manifestations of dengue fever. This part contains 9 questions.

Data collection technique:

Researcher has be visits the selected primary health care setting after getting the approval from the ministry of health . The researcher has be obtained permission from primary health care setting director and participants.

After the arrival of the participants has be explained the purpose of the study to all participants attending .

Data entry and analysis:

The Statistical Package for Social Sciences (SPSS) software version 24.0 has be used for data entry and analysis. Descriptive statistics (e.g., number, percentage) and analytic statistics using Chi-Square tests (χ^2) to test for the association and the difference between two categorical variables were applied. A p-value ≤ 0.05 will be considered statistically significant.

Pilot study

A pilot study has be conducted in primary health care patient's the same sector due to the similarity to the target group using the same questionnaire to test the methodology of the study. As a feedback, the questionnaire will be clear and no defect has be detected in the methodology

Ethical considerations

Permission from the Makkah joint program Family Medicine program has be obtained. Permission from the Directorate of health , verbal consents from all participants in the questionnaire were obtained. All information was kept confidential, and results have be submitted to the department as feedback .

Budget: Self-funded

Result

Table 1. Distribution of the demographic characteristics of about symptoms and sign of the dengue fever in the participants . (n=200)

	N	%
Age		
<30years	46	23
30-45 years	68	34
45-60 years	54	27
More than 60 years	32	16
Sex		
Female	84	42

¹Received 10 August 2023; Accepted 25 August 2023

Male	116	58
Experience		
<2 y	86	43
2-10 y	70	35
>10 y	44	22
Specialty		
Internal medicine	78	39
Surgery	24	12
Family medicine	38	19
Pediatrics	56	28
Ear, nose, throat	4	2
Marital status		
Single	68	34
Married	98	49
Widow	20	10
Divorced	14	7
Reported a dengue case		
Yes	150	75
No	50	25

This table 1 shows that the majority of participant were (34.0%) were aged from 30-45 years of age, while 45-60 of participant were (27.0%) followed by aged < 30 years (23.0%), regarding sex, more than half of participant (58.0%) were male while female were (42.0%) regarding experience the majority of participant <2 y were (43.0%) while 2-10 y were (35.0%) but the > 10 y were (22.0%) , regarding specialty the majority of participant internal medicine were (39 .0%) while pediatrics were (28.0%) but family medicine were (19.0%), regarding marital status, the majority of participant married were (49.0%) while single were(34.0%) but widow were (10.0%), regarding reported a dengue case more than half of participant answer Yes were (75.0%) but answer No were (25.0%) .

Table 2. Distribution of the knowledge about symptoms of the dengue fever in the participants

	Physician				Chi square	
	Yes		No		X²	P-value
	N	%	N	%		
Many people experience no signs or symptoms of a dengue infection DF?	136	68	64	32	25.920	0.000

¹Received 10 August 2023; Accepted 25 August 2023

When symptoms do occur, they may be mistaken for other illnesses Such as the flu DF?	50	25	150	75	50.000	0.000
Usually symptoms begin four to 10 days after you are bitten by an infected mosquito DF?	62	31	138	69	28.880	0.000
Dengue fever causes a high fever — 104 F (40 C) — DF?	24	12	176	88	115.520	0.000
Most people recover within a week or so DF?	38	19	162	81	76.880	0.000
In some cases, symptoms worsen and can become life-threatening called severe DF?	70	35	130	65	18.000	0.000
Do you know Knowledge about symptoms dengue hemorrhagic fever or dengue shock syndrome .	44	22	156	78	62.720	0.000
A symptom of severe dengue happens when your blood vessels become damaged and leaky. And the number of clot-forming cells (platelets) in your bloodstream drops .	48	24	152	76	54.080	0.000
Symptoms of DF this can lead to shock, internal bleeding, organ failure and even death	68	34	132	66	20.480	0.000
A symptom of dengue fever damage to lymph and blood vessels, bleeding from the nose and gums, enlargement of the liver	44	22	156	78	62.720	0.000

Regarding distribution of the knowledge about symptoms of the dengue fever in the participants show regarding the many people experience no signs or symptoms of a dengue infection DF the majority of participant a statistical significant while P-value= 0.000 and X^2 25.920 increased in answer Yes were (68.0%) while answer No were (32.0%), regarding when symptoms do occur, they may be mistaken for other illnesses Such as the flu DF the majority of participant a statistical significant while P-value= 0.000 and X^2 50.000 increased in answer No were (75.0%) while answer Yes were (25.0%), regarding usually symptoms begin four to 10 days after you are bitten by an infected mosquito DF the majority of participant a statistical significant while P-value= 0.000 and X^2 28.880 increased in answer No were (69.0%) while answer Yes were (31.0%), regarding dengue fever causes a high fever — 104 F (40 C) — DF the majority of participant a statistical significant while P-value= 0.000 and X^2 115.520 increased in answer No were (88.0%) while answer Yes were (12.0%), regarding Most people recover within a week or so DF the majority of participant a

¹Received 10 August 2023; Accepted 25 August 2023

statistical significant while P-value= 0.000 and X^2 76.880 increased in answer No were (81.0%) while answer Yes were (19.0%), regarding some cases, symptoms worsen and can become life-threatening called severe DF the majority of participant a statistical significant while P-value= 0.000 and X^2 18.000 increased in answer No were (65.0%) while answer Yes were (35.0%), regarding you know Knowledge about symptoms dengue hemorrhagic fever or dengue shock syndrome the majority of participant a statistical significant while P-value= 0.000 and X^2 62.720 increased in answer No were (78.0%) while answer Yes were (22.0%) , regarding A symptom of severe dengue happens when your blood vessels become damaged and leaky, the number of clot-forming cells (platelets) in your bloodstream drops the majority of participant a statistical significant while P-value= 0.000 and X^2 54.080 increased in answer No were (76.0%) while answer Yes were (24.0%) , regarding symptoms of DF this can lead to shock, internal bleeding, organ failure and even death the majority of participant a statistical significant while P-value= 0.000 and X^2 20.480 increased in answer No were (34.0%) while answer Yes were (66.0%), regarding A symptom of dengue fever damage to lymph and blood vessels, bleeding from the nose and gums, enlargement of the liver the majority of participant a statistical significant while P-value= 0.000 and X^2 62.720 increased in answer No were (78.0%) while answer Yes were (22.0%) .

Table 3. Distribution of the knowledge about sign of the dengue fever in the participants

	Yes		No		Chi-square	
	N	%	N	%	X^2	P-value
1. The warning symptom usually begin the first day or two after your fever goes away	114	57	86	43	3.920	0.048
2. Is fever a symptom of DF?	130	65	70	35	18.000	0.000
3. Is headache a symptom of DF?	110	55	90	45	2.000	0.157
4. Is joint pain a symptom of DF?	134	67	66	33	23.120	0.000
5. Is muscle pain a symptom of DF?	130	65	70	35	18.000	0.000
6. Is pain behind the eyes a symptom of DF?	102	51	98	49	0.080	0.777
7. Are nausea/vomiting symptoms of DF?	134	67	66	33	23.120	0.000
8. Is rash a symptom of DF?	144	72	56	28	38.720	0.000
9. Is diarrhea common in DF?	80	40	120	60	8.000	0.005
10. Is stomach pain common in DF?	98	49	102	51	0.080	0.777
11. Is Swollen glands a symptom of DF	74	37	126	63	13.520	0.000

Regarding distribution of the knowledge about sign of the dengue fever in the participants show regarding the warning symptom usually begin the first day or two after your fever goes

¹Received 10 August 2023; Accepted 25 August 2023

away the majority of participant a statistical significant while P-value= 0.048 and X^2 3.920 increased in answer Yes were (57.0%) while answer No were (43.0%), regarding fever a symptom of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 18.000 increased in answer Yes were (65.0%) while answer No were (35.0%), regarding headache a symptom of DF the majority of participant no statistical significant while P-value= 0.157 and X^2 2.000 increased in answer Yes were (55.0%) while answer No were (45%), regarding joint pain a symptom of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 23.120 increased in answer Yes were (67.0%) while answer No were (33.0%), regarding muscle pain a symptom of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 18.000 increased in answer Yes were (65.0%) while answer No were (35.0%), regarding pain behind the eyes a symptom of DF the majority of participant no statistical significant while P-value= 0.777 and X^2 0.080 increased in answer Yes were (51.0%) while answer No were (49.0%), regarding nausea/vomiting symptoms of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 32.120 increased in answer Yes were (67.0%) while answer Yes were (33.0%) , regarding is rash a symptom of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 38.720 increased in answer Yes were (72.0%) while answer No were (28.0%) , regarding is diarrhea common in DF the majority of participant a statistical significant while P-value= 0.005 and X^2 8.000 increased in answer No were (60.0%) while answer Yes were (40.0%), regarding is stomach pain common in DF the majority of participant no statistical significant while P-value= 0.777 and X^2 13.520 increased in answer No were (63.0%) while answer Yes were (37.0%) , regarding is Swollen glands a symptom of DF the majority of participant a statistical significant while P-value= 0.000 and X^2 13.520 increased in answer No were (63.0%) while answer Yes were (37.0%) .

Table (4) Distribution of the knowledge of practitioner's among the dengue fever

	Knowledge		Chi-square	
	N	%	X^2	P-value
Weak	30	15	33.880	<0.001*
Average	74	37		
High	96	48		
Total	200	100		

Regarding Distribution of the knowledge of practitioner's among the dengue fever show the majority of participant a statistical significant while P-value= 0.001 and X^2 33.880 increased in high knowledge were (48.0%) followed by average were (37.0%) while weak were (15.0%) the total were (100.0%) .

Figure (1) Distribution of the knowledge of practitioner's among the dengue fever

¹Received 10 August 2023; Accepted 25 August 2023




Table 4: Distribution of the Practice of physicians regarding dengue fever

	N	%
How many suspected dengue cases do you see/week?		
Zero	74	37
1-5	90	45
6-10	36	18
Do you look for skin or mucous membrane manifestation if you suspect dengue?		
Always	150	75
Most of the time	20	10
Sometimes	22	11
Never	8	4
Do you perform blood pressure in any suspected case of dengue fever?		
Always	132	66
Most of the time	38	19
Sometimes	20	10
Never	10	5
Do you perform a dengue test if you suspect dengue?		
Always	62	31
Most of the time	36	18
Sometimes	90	45
Never	12	6

¹Received 10 August 2023; Accepted 25 August 2023

If you responded “always” or “most of the time” to above question, which of the following dengue diagnostic tests do you utilize most frequently?		
Leukocyte, platelets, and hematocrit	68	34
Dengue serology (IgM/IgG)	80	40
Dengue reverse-transcription polymerase chain reaction (RT-PCR)	10	5
Viral culture	24	12
	18	9
If you responded “sometimes” or “never” to first question, please explain briefly why you usually do not perform a dengue diagnostic test if you suspect dengue		
CBC is enough	178	89
Test not available	22	11
Please indicate the platelet count for which you would normally refer patients for hospital management.		
<100,000/mm ³	14	7
<80,000/mm ³	20	10
<50,000/mm ³	2	1
Platelet count is not an indicator	164	82

This table 4 regarding distribution of the Practice of physicians regarding dengue fever show regarding the many suspected dengue cases do you see/week the majority of participant 1-5 were (45.0%) while zero were (37.0%) followed by 6-10 were (18.0%), regarding you look for skin or mucous membrane manifestation if you suspect dengue more than half of participant answer always were (75.0%) while sometimes were (11.0%) but most of the time were (10.0%), regarding you perform blood pressure in any suspected case of dengue fever the majority of participant answer always were (66.0%) while most of the time were (19.0%) but sometimes were (10.0%) , regarding you perform a dengue test if you suspect dengue the majority of participant sometimes were (45.0%) while always were (31.0%) but most of the time were (18.0%), regarding you responded “always” or “most of the time” to above question, which of the following dengue diagnostic tests do you utilize most frequently the majority of participant dengue serology (IgM/IgG) were (40.0%) while Leukocyte, platelets, and hematocrit were(34.0%) but chain reaction (RT-PCR) were (12.0%), regarding you responded “sometimes” or “never” to first question, please explain briefly why you usually do not perform a dengue diagnostic test if you suspect dengue more than half of participant CBC is enough were (89.0%) but Test not available were (11.0%), regarding indicate the platelet count for which you would normally refer patients for hospital management more than half of participant Platelet count is not an indicator were (82.0%) but <80,000/mm³ were (10.0%)

Table 4 continua: Distribution of the Practice of physicians regarding dengue fever

	N	%
How frequently do you advice oral fluids and rest to dengue patients?		

¹Received 10 August 2023; Accepted 25 August 2023

Always	16	8
Most of the time	22	11
Sometimes/never	162	81
How frequently do you prescribe paracetamol to dengue patients?		
Always	42	21
Most of the time	118	59
Sometimes/never	40	20
How frequently do you perform daily full blood counts to monitor patients with suspected or confirmed dengue?		
Always	32	16
Most of the time	106	53
Sometimes	24	12
Never	38	19
How frequently do you advice dengue patient to avoid mosquito bite?		
Always	132	66
Most of the time	48	24
Sometimes/never	20	10
Does your center provide to the patient with any type of preventive method (repellent, nets, etc.) to avoid mosquito bite?		
Always	24	12
Most of the time	148	74
Sometimes	24	12
Never	4	2

This table 4 continua regarding frequently do you advice oral fluids and rest to dengue patients the majority of participant sometimes/never were (81.0%) while most of the time were (11.0%) followed by always were (8.0%), regarding frequently do you prescribe paracetamol to dengue patients more than half of participant answer most of the time were (59.0%) while always were (21.0%) but sometimes/never were (20.0%), regarding frequently do you perform daily full blood counts to monitor patients with suspected or confirmed dengue the majority of participant answer most of the time were (53.0%) while never were (19.0%) but always were (16.0%) but sometimes were (12.0%), regarding frequently do you advice dengue patient to avoid mosquito bite the majority of participant always were (66.0%) while most of the time were (24.0%) but sometimes were (10.0%), regarding your center provide to the patient with any type of preventive method (repellent, nets, etc.) to avoid mosquito bite the majority of participant most of the time were (74.0%) while always and sometimes were(12.0%) but never were (2.0%),

Table (5) Distribution of the Practices of practitioner's among the dengue fever


	Practices		Chi-square	
			X ²	P-value
Weak	44	22	21.280	<0.001*

¹Received 10 August 2023; Accepted 25 August 2023

Average	96	48		
High	60	30		
Total	200	100		

Regarding Distribution of the practices of practitioner's among the dengue fever show the majority of participant a statistical significant while P-value= 0.001 and X^2 21.280 increased in average of practices were (48.0%) followed by high were (30.0%) while weak were (22.0%) the total were (100.0%) .

Figure (2) Distribution of the Practice of practitioner's among the dengue fever



Discussion

Dengue fever is the most prevalent vector-borne viral disease in the twenty-first century. It can be found in rural regions, but it is more prevalent in metropolitan and peri-urban areas. In our study assessment of the Knowledge and practices of dengue fever symptoms among health care workers at the primary healthcare level in Makkah City at Saudi Arabia 2023. This study reveals that the majority of participant were (34.0%) were aged from 30-45 years of age, sex, more than half of participant (58.0%) were male, experience the majority of participant <2 y were (43.0%), specialty the majority of participant internal medicine were (39.0%), majority of participant married were (49.0%) while single were (34.0%) but widow were (10.0%), regarding reported a dengue case more than half of participant answer Yes were (75.0%). This results is in accordance with Collij, et al .2021 who found that the majority of study participants were male, married (25). A previous systematic review showed the wide variation of the definitions and cutoff values used by

¹Received 10 August 2023; Accepted 25 August 2023

researchers to classify dengue patients (11). Likewise, the current study found a wide variation in defining the warning signs among physicians in clinical settings. Of these seven warning signs, the HCPs have not agreed on any signs, even with the definition of liver enlargement, which was predefined in WHO 2009 classification. This variation may be linked to the local modification of the guidelines, adoption of national guidelines, lack of proper training, and variation in clinical practice in different regions. (22)

Dengue fever is a major parvovirus-borne infectious disease in tropical and subtropical regions of the world. The disease is generally mild and self-limited, but some patients may develop a severe form of infection such as dengue hemorrhagic fever/dengue shock syndrome, which often leads to death. Currently, there is no vaccine available against dengue virus. The major determinants of dengue control are vector eradication, early case recognition, and adequate clinical management (25,26)

The results of the present study showed the participants have a high and high level of knowledge about signs, symptoms of dengue fever. also statistical significant correlation between knowledge symptoms of the dengue fever in the participants (see table 2,3), regarding the distribution of the knowledge of practitioner's among the dengue fever in our study reported that majority of participant a statistical significant while P-value= 0.001 and χ^2 33.880 increased in high knowledge were (48.0%) followed by average were (37.0%) while weak were (15.0%) the total were (100.0%) (See table 4) in similar study men made up more than half of the participants in the present research . This is because guys outnumber girls in Yemeni university study. This result was unanimous (9). Another possibility could be physicians having more contact with patients (21)

results is in opposite with Ho et al, 2013 who conduct a study to assess Knowledge, attitude, and practice of dengue disease among healthcare professionals in southern Taiwan and found a lack of knowledge about important clinical characteristics of dengue (31). Previous study of the knowledge about dengue fever in the primary health-care medical practitioner's in KSA concerning control of dengue virus showed the lack of knowledge about sign and symptoms as the most common problem.(33,35). Another study similar found almost a third of PHC physicians had insufficient knowledge about important investigations of dengue as well as prevention measures toward DF. In southern Taiwan (35)

However, the study results was in the same line with pan et al 2020 who found that Knowledge of DF as reported by the respondents was average, respondents' knowledge of DF prevention and protection was moderate, attitude of DF as reported by the respondents was at good level, practice of DF prevention as reported by the respondents was at high level and there was statistical significant between the knowledge and attitude of the respondents and practice of DF prevention (36)

In addition, the results of the study was far away Alhaeli, et al. 2016 who conduct a study to assess Knowledge, Attitude and Practice Regarding Dengue Fever among the Healthy Population of Highland and Lowland Communities in Central Nepal. They reveal a low, unsatisfactory level of knowledge about the disease. While being in line with our study finding in relation to presence of significantly positive correlation among knowledge and practice. Among the socio-demographic variables, the education level of the participants, age, sex , marital status and income were an independent predictors of knowledge level. Education

¹Received 10 August 2023; Accepted 25 August 2023

level and interaction between the marital status, income and age group of the participants were independent predictors of practice level (28,30).

similar our study shows that also regarding distribution of the practices of practitioner's among the dengue fever show the majority of participant a statistical significant while P-value= 0.001 and X^2 21.280 increased in average of practices were (48.0%) followed by high were (30.0%) while weak were (22.0%) the total were (100.0%) (see table 4,5)

Conclusion

Best first line early management for SARS CoV-2 and dengue virus and/or other described an yet to be explored respiratory and GIT viral (co)infections, saving precious lives especially in the developed countries that were and are the most affected countries as regards to dengue fever mortality and relieving the health care professionals as well as the health care authorities globally and especially in the developing countries This study showed an average of Knowledge about symptoms and sign of dengue fever among HCPs in KSA. Future continued medical/nursing education should place more emphasis on these factors to improve dengue control in this demographic area, also the study we conclude that dengue fever is prevalent in Saudi Arabia. Thus, strong and effective health education programs regarding dengue fever about symptoms, sign and risk factors are recommended to help prevent dengue fever . However, the paucity of large epidemiological studies limits generalizability of such evidence. Future studies in Saudi Arabia should focus on the expansion of dengue fever to other cities in the Kingdom. Larger epidemiological studies are needed for estimating the true burden and incidence of dengue fever in the Saudi population. Currently, there are few epidemiological studies about dengue fever.

Reference

- [1] Kok, B. H., Lim, H. T., Lim, C. P., Lai, N. S., Leow, C. Y., & Leow, C. H. (2022). Dengue virus infection—a review of pathogenesis, vaccines, diagnosis and therapy. *Virus research*, 199018.
- [2] Wang, X., Liu, Y., Wang, H., Sun, Q., Cheng, P., Zhang, Y., ... & Liu, H. (2023). Wolbachia and Mosquitoes: Exploring Transmission Modes and Coevolutionary Dynamics in Shandong Province, China.
- [3] Poterek, M. L., Vogels, C. B., Grubaugh, N. D., Ebel, G. D., Alex Perkins, T., & Cavany, S. M. (2022). Interactions between seasonal temperature variation and temporal synchrony drive increased arbovirus co-infection incidence. *Royal Society Open Science*, 9(10), 220829.
- [4] Kelleni, M. (2023). Kelleni's Protocol including Nitazoxanide and NSAIDs to Manage Dengue Virus Disease: A Novel Potential Game-Changer.
- [5] Brereton, P. (2022). Essential Concepts of Environmental Communication: An A-Z Guide.
- [6] Dafalla, O., Abdulhaq, A. A., Almutairi, H., Noureldin, E., Ghzwani, J., Mashi, O., ... & Eisa, Z. M. (2023). The emergence of an imported variant of dengue virus serotype 2 in the Jazan region, southwestern Saudi Arabia. *Tropical diseases, travel medicine and vaccines*, 9(1), 1-9.

¹Received 10 August 2023; Accepted 25 August 2023

- [7] Alshabi, A., Marwan, A., Fatima, N., Madkhali, A. M., Alnagai, F., Alhazmi, A., ... & Refaei, T. (2022). Epidemiological screening and serotyping analysis of dengue fever in the Southwestern region of Saudi Arabia. *Saudi Journal of Biological Sciences*, 29(1), 204-210.
- [8] Mohammed Yusuf, A., & Abdurashid Ibrahim, N. (2019). Knowledge, attitude and practice towards dengue fever prevention and associated factors among public health sector health-care professionals: in Dire Dawa, eastern Ethiopia. *Risk Management and Healthcare Policy*, 91-104.
- [9] Giang, H. T. N., Sayed, A. M., Dang, T., Iqtadar, S., Tuan, N. M., Khiem, N. T., ... & Huy, N. T. (2021). Survey of knowledge, attitude and practice of healthcare professionals on dengue transmission, diagnosis and clinical classification. *BMC infectious diseases*, 21, 1-11.
- [10] Dalpadado, R., Amarasinghe, D., Gunathilaka, N., & Ariyaratna, N. (2022). Bionomic aspects of dengue vectors *Aedes aegypti* and *Aedes albopictus* at domestic settings in urban, suburban and rural areas in Gampaha District, Western Province of Sri Lanka. *Parasites & Vectors*, 15(1), 1-14.
- [11] Egid, B. R., Coulibaly, M., Dadzie, S. K., Kamgang, B., McCall, P. J., Sedda, L., ... & Wilson, A. L. (2022). Review of the ecology and behaviour of *Aedes aegypti* and *Aedes albopictus* in Western Africa and implications for vector control. *Current research in parasitology & vector-borne diseases*, 2, 100074.
- [12] Kong, L., Xiao, J., Yang, L., Sui, Y., Wang, D., Chen, S., ... & Gu, J. (2023). Mosquito densovirus significantly reduces the vector susceptibility to dengue virus serotype 2 in *Aedes albopictus* mosquitoes (Diptera: Culicidae). *Infectious Diseases of Poverty*, 12(1), 1-14.
- [13] Huynh, T. T. T., & Minakawa, N. (2022). A comparative study of dengue virus vectors in major parks and adjacent residential areas in Ho Chi Minh City, Vietnam. *PLOS Neglected Tropical Diseases*, 16(1), e0010119.
- [14] Badolo, A., Sombié, A., Yaméogo, F., Wangrawa, D. W., Sanon, A., Pignatelli, P. M., ... & McCall, P. J. (2022). First comprehensive analysis of *Aedes aegypti* bionomics during an arbovirus outbreak in west Africa: Dengue in Ouagadougou, Burkina Faso, 2016–2017. *PLOS Neglected Tropical Diseases*, 16(7), e0010059.
- [15] Liu, S. Y., Chien, T. W., Yang, T. Y., Yeh, Y. T., Chou, W., & Chow, J. C. (2021). A bibliometric analysis on dengue outbreaks in tropical and sub-tropical climates worldwide since 1950. *International Journal of Environmental Research and Public Health*, 18(6), 3197.
- [16] Wong, J. M., Adams, L. E., Durbin, A. P., Muñoz-Jordán, J. L., Poehling, K. A., Sánchez-González, L. M., ... & Paz-Bailey, G. (2022). Dengue: a growing problem with new interventions. *Pediatrics*, 149(6), e2021055522.
- [17] Khan, M. B., Yang, Z. S., Lin, C. Y., Hsu, M. C., Urbina, A. N., Assavalapsakul, W., ... & Wang, S. F. (2023). Dengue overview: An updated systemic review. *Journal of Infection and Public Health*.
- [18] Macedo, G. A., Gonin, M. L. C., Pone, S. M., Cruz, O. G., Nobre, F. F., & Brasil, P. (2014). Sensitivity and specificity of the World Health Organization dengue

¹Received 10 August 2023; Accepted 25 August 2023

- classification schemes for severe dengue assessment in children in Rio de Janeiro. *PloS one*, 9(4), e96314.
- [19] Horstick, O., Farrar, J., Lum, L., Martinez, E., San Martin, J. L., Ehrenberg, J., ... & Kroeger, A. (2012). Reviewing the development, evidence base, and application of the revised dengue case classification. *Pathogens and global health*, 106(2), 94-101.
- [20] Johnson, A. K. (2016). *Surveillance of Sexually Transmitted Infections: Exploring the Utility of Internet Search Engine Data* (Doctoral dissertation, University of Illinois at Chicago).
- [21] Low, G. K. K., Ogston, S. A., Yong, M. H., Gan, S. C., & Chee, H. Y. (2018). Global dengue death before and after the new World Health Organization 2009 case classification: A systematic review and meta-regression analysis. *Acta tropica*, 182, 237-245.
- [22] Peraza, O. C., Torres, E. M., Rodr guez, A. M., & Tirado, M. G. G. (2022). Usefulness of warning signs in the management of dengue patients. *Revista Cubana de Medicina Tropical*, 74(2).
- [23] Huits, R., Angelo, K. M., Amatya, B., Barkati, S., Barnett, E. D., Bottieau, E., ... & Schwartz, E. (2023). Clinical Characteristics and Outcomes Among Travelers With Severe Dengue: A GeoSentinel Analysis. *Annals of Internal Medicine*.
- [24] Junior, J. B. S., Massad, E., Lobao-Neto, A., Kastner, R., Oliver, L., & Gallagher, E. (2022). Epidemiology and costs of dengue in Brazil: a systematic literature review. *International Journal of Infectious Diseases*.
- [25] Collij, L. E., Salvadó, G., Shekari, M., Lopes Alves, I., Reimand, J., Wink, A. M., ... & AMYPAD consortium. (2021). Visual assessment of [18 F] flutemetamol PET images can detect early amyloid pathology and grade its extent. *European Journal of Nuclear Medicine and Molecular Imaging*, 48, 2169-2182.
- [26] van Gennip, L., Haverkamp, F. J., Sir,  ., & Tan, E. C. (2022). A multicentre cross-sectional survey study on acute wound classification in the emergency department and its interobserver variability. *Scientific Reports*, 12(1), 9901.
- [27] Htun, T. P., Xiong, Z., & Pang, J. (2021). Clinical signs and symptoms associated with WHO severe dengue classification: a systematic review and meta-analysis. *Emerging Microbes & Infections*, 10(1), 1116-1128.
- [28] Alhaeli, A., Bahkali, S., Ali, A., Househ, M. S., & El-Metwally, A. A. (2016). The epidemiology of Dengue fever in Saudi Arabia: A systematic review. *Journal of infection and public health*, 9(2), 117-124.
- [29] Altassan, K. K., Morin, C., Shocket, M. S., Ebi, K., & Hess, J. (2019). Dengue fever in Saudi Arabia: A review of environmental and population factors impacting emergence and spread. *Travel medicine and infectious disease*, 30, 46-53.
- [30] Sahu, M. K., & Tiwari, S. P. (2023). A Systemic Review on Recent Outbreaks of Dengue and Newer US FDA Approved Drug. *J Infec Dise and Vir Res 2 (2)*, 01, 6.
- [31] Unissa, R., Jyothirmayi, B., Mounica, A., Tejaswini, L., Latha, G. A., Nishitha, G., & Swati, C. (2018). Dengue. *Asian Journal of Research in Pharmaceutical Science*, 8(4), 192-202.
- [32] Nanyingi, M. O. (2018). *Spatial epidemiology and predictive modelling of Rift valley fever in Garissa county, Kenya* (Doctoral dissertation, University of Nairobi).

¹Received 10 August 2023; Accepted 25 August 2023

- [33] Dsouza, R. P., Rodrigues, D. E., & Saldanha, P. M. (2022). Effectiveness of school-based video-assisted health education program on mosquito-borne disease among upper primary children. *Journal of Health and Allied Sciences NU*, 13(01), 098-102.
- [34] Tsheten, T., Clements, A. C., Gray, D. J., Gyeltshen, K., & Wangdi, K. (2021). Medical practitioner's knowledge on dengue management and clinical practices in Bhutan. *PloS one*, 16(7), e0254369.
- [35] Tsheten, T., Clements, A. C., Gray, D. J., Gyeltshen, K., & Wangdi, K. (2021). Medical practitioner's knowledge on dengue management and clinical practices in Bhutan. *PloS one*, 16(7), e0254369.
- [36] Pan, C. Y., Liu, W. L., Su, M. P., Chang, T. P., Ho, H. P., Shu, P. Y., ... & Chen, C. H. (2020). Epidemiological analysis of the Kaohsiung city strategy for dengue fever quarantine and epidemic prevention. *BMC Infectious Diseases*, 20(1), 1-9.