

Active Monitoring of Person Exposed to Patients with Confirmed COVID - 19

Vignesh P

Department of General Pathology,
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
Chennai -77,
Tamilnadu.
Mail id : 151701042.sdc@saveetha.com

Dr. Brundha M.P .

Associate professor,

Department of General Pathology
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences No: 162, PH Road
Chennai - TamilNadu
India - 600077
Email: brundha.sdc@saveetha.com

Corresponding Author

Dr. Brundha M.P

Associate professor,

Department of General Pathology
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical Sciences,
Chennai - TamilNadu
India - 600077
Email: brundha.sdc@saveetha.com

ABSTRACT

Coronavirus disease 2019 is an illness brought about by extreme severe respiratory condition coronavirus (SARS-CoV-2). It was first determined in the 2019 December month in Wuhan fish market in China, and has brought about a progressing epidemic. The first case might be recorded back on 17 November 2019.

Symptoms of the COVID-19 include cough, fever, dyspnea, fatigue, and ability to smell and taste is lost. While a large portion of the cases show mild signs and symptoms, some of them arrive at acute respiratory distress syndrome (ARDS) likely encouraged by a cytokine storm,

septic stun, multi-organ failure , and blood clumps. After exposure the time it takes to begin symptoms is generally around six days or it may go from two to fourteen days.

The virus is initially spread between person to person during close contact, most commonly by means of small droplets that come out by sneezing, talking, and coughing. The droplets usually fall onto things and not travel through air over long distances. Less ordinarily, individuals may get infected by contacting a contaminated surface at that point then touching their face. It's generally infectious toward the start of the initial three days after the beginning of indications, despite the fact that the spread is achievable before manifestations show up, and from individuals that don't show side effects.

Keywords: COVID- 19, Transmissions, Management, Global Pandemic, Prevention.

INTRODUCTION

In 1930, the first coronavirus was discovered. An acute respiratory infection of tamed Chicken was caused by injections of bronchitis Virus (IBV)(Estola, 1970), Anther Schalk, and M.C. Hawn described in 1931 a brand new respiratory virus of Chicken in North Dakota (Fabricant, 1998). Later, Between 1930 - 1940 three more different animal corona Viruses were Cultivated and were kept isolated, at that period they did not realize that these various different viruses were related (McIntosh, 1974). In 1960 the first human coronavirus was discovered. The isolation was done using two different methods in the United nation and in America. In 1960 same year Original research was done about Virus isolation from Common cold occurring in a residential School. From the general population the schoolboy was tested. Malcolm Byone, E.C. Kendall, and David Tyrrell working at the Common Cold Unit of the British Medical Research Council in 1960 isolated from a boy a novel common cold virus B814. (Kendall, Bynoe and Tyrrell, 1962; Kahn and McIntosh, 2005) The discovered virus was not technically cultivated using normal standard methods which had successfully cultivated rhinoviruses, adenoviruses and other common cold viruses. Like this furthermore, viruses are cultivated and monitored (Corman *et al.*, 2014) After a few decades, the coronavirus strain B814 was lost. (Su *et al.*, 2016; Zhu *et al.*, 2020)Our institution is passionate about high quality evidence based research and has excelled in various fields ((Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; Vijayashree Priyadharsini, Smiline Girija and Paramasivam, 2018; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Ramadurai *et al.*, 2019; Sridharan *et al.*, 2019; Vijayashree Priyadharsini, 2019; Chandrasekar *et al.*, 2020; Mathew *et al.*, 2020; R *et al.*, 2020; Samuel, 2021)

Infected individuals are able to contaminate the environment. The interaction of the complementary cell receptor and coronavirus spike protein is central in determining the infectivity, tissue tropism and species range of the released virus. (Masters, 2006) Coronavirus target epithelial cells. (Cui, Li and Shi, 2019) They are transmitted from one individual to another individual depending on the corona Virus species by either aerosol fomite or orofecal

route (Decaro, 2011). Patients infected with coronavirus Can present a wide range of signs and symptoms extending from mild to severe. cough, Fever, and dyspnea are the most common symptoms reported, and 31% of cases. (Li *et al.*, 2020) Patients that develop acute respiratory distress Syndrome may worsen rapidly and die of multiple organ failure. (Wang *et al.*, 2020). Our team has rich experience in research and we have collaborated with numerous authors over various topics in the past decade (Ariga *et al.*, 2018; Basha, Ganapathy and Venugopalan, 2018; Hannah *et al.*, 2018; Hussainy *et al.*, 2018; Jeevanandan and Govindaraju, 2018; Kannan and Venugopalan, 2018; Kumar and Antony, 2018; Manohar and Sharma, 2018; Menon *et al.*, 2018; Nandakumar and Nasim, 2018; Nandhini, Babu and Mohanraj, 2018; Ravintha and Jayalakshmi, 2018; Seppan *et al.*, 2018; Teja, Ramesh and Priya, 2018; Duraisamy *et al.*, 2019; Gheena and Ezhilarasan, 2019; Hema Shree *et al.*, 2019; Rajakeerthi and Ms, 2019; Rajendran *et al.*, 2019; Sekar *et al.*, 2019; Sharma *et al.*, 2019; Siddique *et al.*, 2019; Janani, Palanivelu and Sandhya, 2020; Johnson *et al.*, 2020; Jose, Ajitha and Subbaiyan, 2020).

COVID - 19 OUTBREAK

An outbreak of viral infection of unknown origin was first determined in Wuhan fish market, Hubei, provinces, china. It was started in December 2019. There Case was epidemiologically connected to the human seafood market. (Almeida and Tyrrell, 1967) Inoculation of bronchoalveolar lavage liquid acquired from patients with viral infection of unknown Origin into human airway epithelial cell and Huh7 and Vero E6 cell lines led to the isolation of a SARS-CoV-2, novel coronavirus, earlier named 2019-nCov.(Balasubramanian, 2020) (Kapikian *et al.*, 1969)

Coronaviruses are positive single-stranded RNA viruses surrounded by an envelope belonging to the family Coronaviridae. They are classified into four genera Alphacoronavirus, Betacoronavirus, Gammacoronavirus, and Deltacoronavirus. So far seven human coronaviruses (HCoVs) have been identified, which fall within the Alpha- and Betacoronavirus genera.

SARS-Cov are animal origin, they cause severe respiratory syndrome and are often fatal. From the beginning of the pandemic in late December 2019, SARS-COV-2 has now spread to all continents as of May 21, 2020, the world health organization. Communicated 5,101,526 Confirmed cases and 329,203 deaths globally. The name Coronavirus was coined by June Almeida and David Tyrrell (He is the one who found novel common cold B814 in a Schoolboy). (Woo *et al.*, 2005) Coronavirus can infect the gastrointestinal and upper respiratory tract of mammals and birds. (Bande *et al.*, 2015)

MODE OF TRANSMISSION OF COVID - 19

Respiratory infections might be transmitted through droplets of different sizes: once the droplet particle unit area is $>5\text{-}10 \mu\text{m}$ in diameter they're called respiratory droplets, and once the unit area $<5\mu\text{m}$ in diameter, they're called as droplet nuclei.(Conly *et al.*, 2011) In step with current proof, the COVID - 19 virus is principally transmitted between people through respiratory droplets and by contacting individuals. (Chan *et al.*, 2020; Liu *et al.*, 2020)

Droplet transmission happens once someone is in close contact (within one meter) with somebody who has symptoms (e.g., coughing or sneezing) and is thus in danger of getting his/her mucosae (mouth and nose) or mucous membrane (eyes) exposed to probably infective metabolism droplets. (Ong *et al.*, 2020)

Droplet transmission is entirely different from airborne transmission since it refers to the presence of microbes inside droplet nuclei, (Burke *et al.*, 2020) that area unit airborne transmission usually thought-about to be particles $<5\mu\text{m}$ in diameter, will remain within the air for long periods of your time and be transmitted to others over distances longer than one meter.(Chan *et al.*, 2020)

With regards to COVID - 19, the transmission system is additionally potential in explicit conditions and settings during which techniques or bolster medicines that produce pressurized canned products territory unit performed; i.e., manual ventilation before brooding, turning the patient to the inclined position, detaching the patient from the ventilator, non-obtrusive positive-pressure ventilation, careful activity, and heart resuscitation(Brundha, no date a).

There is some proof that COVID - 19 infection could cause internal organ infection and be present in feces. However, up to now, just one study has civilized the COVID - 19 virus from one stool specimen. (Zhang *et al.*, 2020)There are no reports of fecal-oral transmission of the COVID - 19 virus up to now.

FOUR STAGES OF GLOBAL PANDEMIC

As indicated by the report, the first stage is when instances of contamination are brought into a country that was not the source of the disease. At that moment, all the country outside of China that began announcing instances of coronavirus reached at Stage 1 of the outbreak as they revealed their first case of infection (Brundha and Nallaswamy, 2019; Brundha, Pathmashri and Sundari, 2019).If a virus is contained after transmission to just one or two countries, it doesn't turn into a pestilence, however on the off chance that it can't be contained during a limited period of time and spreads over the world, as COVID - 19 did,(Timothy, Samyuktha and Brundha, 2019) it turns into an epidemic.(Saif, no date).

The second stage of a pandemic is when there are instances of local transmission inside the country; the report clarifies (Prashaanthi and Brundha, 2018). This means the individual from

whom the contamination spread to an alternate individual is from the same country. During this stage, the direction of the infection is generally recognized from the source to all the contaminated people. (Kumar and Brundha, 2016).

Community transmission is the third stage of the pandemic, in this stage, predictable with the report; it turns out to be difficult to follow the chain of transmission of the infection in a sizable measure of cases. This recommends the infection has begun circling inside the network and may likewise contaminate those people who neither have headed out to a country experiencing the pandemic nor have come into the contact of an individual contaminated by the infection. During this event, (Hannah *et al.*, 2019) a lockdown turns out to be profoundly significant as an individual can spread the infection, paying little heed to their movement history or the individuals they need are accessible in contact with (Preethikaa and Brundha, 2018).

The fourth phase is when the virus infection gets endemic in certain continues reemerging around the year. This is frequently a phase that the Indian government has mulled over in its choice to handle the infection, the IE report states(Shreya and Brundha, 2017).

These stages of an episode stay uniform over the planet to shape coordination and understanding less difficult, all together that countries are frequently arranged in like manner. Such an order makes it simpler for different countries to force quantifiers that they think will profit, (Kalaiselvi and Brundha, 2016) like India forcing travel limitations on China very right on time inside the epidemic. This was on the grounds that, the report states, around then of your time all the instances of coronavirus in different nations were being imported from China (Harsha and Brundha, 2017).

MANAGEMENT OF PATIENT

The incubation time for coronavirus is said to be extended to 14 days, within a time of 4-5 days from exposure to symptoms onset. One research study revealed that 97% of persons with confirmed COVID - 19 develop symptoms can therefore at intervals of 11.5 days of coronavirus infection(Guan *et al.*, 2020).

The signs and symptoms of novel corona virus present at unwellness onset vary, however over the course of the sickness, (Chen *et al.*, 2020) most persons with COVID - 19 can have Fever, Cough, Anorexia, Fatigue, Sputum production, dyspnea, Myalgia and Atypical symptoms are mostly seen in older adults and persons with past medical history and comorbidities may have delayed presentation of fever and respiratory signs and symptoms. An article concluded, it is referenced around 1,000 hospitalized patients, fever was present in just 44% at hospital admission however later developed in 89% in the hospital. (Huang *et al.*, 2020)confusion, Headache, rhinorrhea, hemoptysis, pharyngitis, vomiting, and diarrhea symptoms are reported less frequent (<10%). (Wang *et al.*, 2020)Dysnomia or ageusia preceding the onset of respiratory symptoms has been anecdotally reported, however additional info is required to grasp its role in characteristic COVID - 19.(Balaji, Brundha and Path, 2016; Ravichandran and Brundha, 2016).

INFECTION PREVENTION AND CONTROL

Immediate recognition, Securing triage, and preventing the source of infection. clinical storage includes a system for assuring all patients at admission, allowing for early identification of possible COVID - 19 and immediate quarantine of patients with Suspected infection in an area isolated from other patients.(Shenoy and Brundha, 2016)Standard precautionary measures include hand and respiratory cleanliness, the utilization of appropriate personal protective equipment (PPE) according to a risk assessment, waste management, environmental cleaning, conventional linens, and sterilization of patient-care supplies. (Adhikari *et al.*, 2020)

In addition to utilizing the standard precautions, all people, including family members and guests, should use contact and droplet precautionary measures before entering the room of suspected or Confirmed COVID -19 patient, Patient should be placed in inadequately ventilated single rooms. For a general wardroom with natural ventilation, sufficient Ventilation is considered (Brundha, no date b) .

60 L/s per patient. Authoritative controls and arrangements for the avoidance and control of transmission of COVID - 19 within the health care setting include, yet may not be constrained to building up practical IPC infrastructure and activities. instructing patients developing policies on early recognition of acute respiratory infection potentially could by Could- 19 infections (Shen *et al.*, 2020).

CONCLUSION

This study is a picture of current research of transmission, management, prevention, and pandemic of ongoing Coronavirus. protect yourself as well as other people around you by knowing the information and taking Appropriate measures, follow the suggestion given by your nearby general wellbeing office.

REFERENCE

1. Adhikari, S. P. *et al.* (2020) 'Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: a scoping review', *Infectious Diseases of Poverty*, 9(1), p. 29. doi: 10.1186/s40249-020-00646-x.
2. Almeida, J. D. and Tyrrell, D. A. (1967) 'The morphology of three previously uncharacterized human respiratory viruses that grow in organ culture', *The Journal of general virology*, 1(2), pp. 175–178. doi: 10.1099/0022-1317-1-2-175.
3. Ariga, P. *et al.* (2018) 'Determination of correlation of width of Maxillary Anterior Teeth using Extraoral and Intraoral Factors in Indian Population: A systematic review', *World journal of dentistry*, 9(1), pp. 68–75. doi: 10.5005/jp-journals-10015-1509.
4. Balaji, S., Brundha, M. P. and Path, D. N. B. (2016) 'Awareness of About Breast Cancer among Dental Surgeons', *Research journal of pharmaceutical, biological and chemical sciences*, 8(8), p. 797. Available at: <https://pdfs.semanticscholar.org/63f4/4173d90b35bffa33eed0aeb52ac547ef1567.pdf>.

5. Balasubramanian, D. M. (2020) *Covid 19 - The New Age Pandemic*. Notion Press. Available at: <https://play.google.com/store/books/details?id=gsrlDwAAQBAJ>.
6. Bande, F. *et al.* (2015) 'Progress and challenges toward the development of vaccines against avian infectious bronchitis', *Journal of immunology research*, 2015, p. 424860. doi: 10.1155/2015/424860.
7. Basha, F. Y. S., Ganapathy, D. and Venugopalan, S. (2018) 'Oral hygiene status among pregnant women', *Journal of advanced pharmaceutical technology & research*, 11(7), p. 3099. doi: 10.5958/0974-360x.2018.00569.3.
8. Brundha, M. P. (no date a) 'A Comparative Study- The Role of Skin and Nerve Biopsy in Hansen's Disease'.
9. Brundha, M. P. (no date b) 'Awareness of Styte'.
10. Brundha, M. P. and Nallaswamy, D. (2019) 'Hide and seek in pathology- A research on game-based histopathology learning', *International Journal of Research in Pharmaceutical Sciences*, 10(2), pp. 1410–1414. doi: 10.26452/ijrps.v10i2.606.
11. Brundha, M. P., Pathmashri, V. P. and Sundari, S. (2019) 'Quantitative Changes of Red Blood cells in Cancer Patients under Palliative Radiotherapy-A Retrospective Study', *Research Journal of Pharmacy and Technology*, 12(2), pp. 687–692. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=2&article=041>.
12. Burke, R. M. *et al.* (2020) 'Active Monitoring of Persons Exposed to Patients with Confirmed COVID-19 - United States, January–February 2020', *MMWR. Morbidity and mortality weekly report*, 69(9), pp. 245–246. doi: 10.15585/mmwr.mm6909e1.
13. Chandrasekar, R. *et al.* (2020) 'Development and validation of a formula for objective assessment of cervical vertebral bone age', *Progress in orthodontics*, 21(1), p. 38. doi: 10.1186/s40510-020-00338-0.
14. Chan, J. F.-W. *et al.* (2020) 'A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster', *The Lancet*, 395(10223), pp. 514–523. doi: 10.1016/S0140-6736(20)30154-9.
15. Chen, N. *et al.* (2020) 'Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study', *The Lancet*, 395(10223), pp. 507–513. doi: 10.1016/S0140-6736(20)30211-7.
16. Conly, J. *et al.* (2011) 'Field evaluation (FE) of the World Health Organization (WHO) interim guidelines (IG) on infection prevention and control (IPC) of epidemic and pandemic-prone acute respiratory diseases (ARD) in health care', *BMC Proceedings*. doi: 10.1186/1753-6561-5-s6-p280.
17. Corman, V. M. *et al.* (2014) 'Antibodies against MERS Coronavirus in Dromedary Camels, Kenya, 1992–2013', *Emerging Infectious Diseases*. doi: 10.3201/eid2008.140596.
18. Cui, J., Li, F. and Shi, Z.-L. (2019) 'Origin and evolution of pathogenic coronaviruses', *Nature Reviews Microbiology*, pp. 181–192. doi: 10.1038/s41579-018-0118-9.
19. Decaro, N. (2011) 'Alphacoronavirus‡: Coronaviridae', *The Springer Index of Viruses*, p. 371. doi: 10.1007/978-0-387-95919-1_56.
20. Duraisamy, R. *et al.* (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295. doi: 10.1097/ID.0000000000000885.

21. Estola, T. (1970) 'Coronaviruses, a new group of animal RNA viruses', *Avian diseases*, 14(2), pp. 330–336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/4316767>.
22. Ezhilarasan, D., Apoorva, V. S. and Ashok Vardhan, N. (2019) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121. doi: 10.1111/jop.12806.
23. Fabricant, J. (1998) 'The early history of infectious bronchitis', *Avian diseases*, 42(4), pp. 648–650. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9876830>.
24. Gheena, S. and Ezhilarasan, D. (2019) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & experimental toxicology*, 38(6), pp. 694–702. doi: 10.1177/0960327119839173.
25. Guan, W.-J. *et al.* (2020) 'Clinical Characteristics of Coronavirus Disease 2019 in China', *The New England journal of medicine*, 382(18), pp. 1708–1720. doi: 10.1056/NEJMoa2002032.
26. Hannah, R. *et al.* (2018) 'Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens', *Journal of advanced pharmaceutical technology & research*, 11(3), p. 1012. doi: 10.5958/0974-360x.2018.00189.0.
27. Hannah, R. *et al.* (2019) 'Liquid Paraffin as a Rehydrant for Air Dried Buccal Smear', *Research Journal of Pharmacy and Technology*, 12(3), pp. 1197–1200. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=3&article=038>.
28. Harsha, L. and Brundha, M. P. (2017) 'Prevalence of dental developmental anomalies among men and women and its psychological effect in a given population', *Research journal of pharmaceutical, biological and chemical sciences*, 9(6), p. 869. Available at: <http://search.proquest.com/openview/1f488cc6e377096f44a87e509aceab79/1?pq-origsite=gscholar&cbl=54977>.
29. Hema Shree, K. *et al.* (2019) 'Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma - a Systematic Review with Meta Analysis', *Pathology oncology research: POR*, 25(2), pp. 447–453. doi: 10.1007/s12253-019-00588-2.
30. Hussainy, S. N. *et al.* (2018) 'Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up', *Journal of conservative dentistry: JCD*, 21(5), pp. 510–515. doi: 10.4103/JCD.JCD_51_18.
31. Janani, K., Palanivelu, A. and Sandhya, R. (2020) 'Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality: an in vivo study', *Brazilian dental science*, 23(1). doi: 10.14295/bds.2020.v23i1.1805.
32. Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 19(4), pp. 273–278. doi: 10.1007/s40368-018-0356-6.
33. Johnson, J. *et al.* (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH',

- Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362. doi: 10.1038/s41440-019-0369-5.
34. Jose, J., Ajitha and Subbaiyan, H. (2020) ‘Different treatment modalities followed by dental practitioners for Ellis class 2 fracture – A questionnaire-based survey’, *The open dentistry journal*, 14(1), pp. 59–65. doi: 10.2174/1874210602014010059.
 35. Kahn, J. S. and McIntosh, K. (2005) ‘History and recent advances in coronavirus discovery’, *The Pediatric infectious disease journal*, 24(11 Suppl), pp. S223–7, discussion S226. doi: 10.1097/01.inf.0000188166.17324.60.
 36. Kalaiselvi, R. and Brundha, M. P. (2016) ‘Prevalence of hysterectomy in South Indian population’, *Research Journal of Pharmacy and Technology*, 9(11), pp. 1941–1944. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=9&issue=11&article=027>.
 37. Kannan, A. and Venugopalan, S. (2018) ‘A systematic review on the effect of use of impregnated retraction cords on gingiva’, *Journal of advanced pharmaceutical technology & research*, 11(5), p. 2121. doi: 10.5958/0974-360x.2018.00393.1.
 38. Kapikian, A. Z. et al. (1969) ‘Isolation from man of “ avian infectious bronchitis virus-like” viruses (coronaviruses) similar to 229E virus, with some epidemiological observations’, *The Journal of infectious diseases*, pp. 282–290. Available at: <https://www.jstor.org/stable/30102315>.
 39. Kendall, E. J. C., Bynoe, M. L. and Tyrrell, D. A. J. (1962) ‘Virus Isolations from Common Colds Occurring in a Residential School’, *BMJ*, pp. 82–86. doi: 10.1136/bmj.2.5297.82.
 40. Kumar, D. and Antony, S. D. P. (2018) ‘Calcified canal and negotiation-A review’, *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3727. doi: 10.5958/0974-360x.2018.00683.2.
 41. Kumar, M. D. and Brundha, M. P. (2016) ‘Awareness about nocturia-A questionnaire survey’, *Research Journal of Pharmacy and Technology*, 9(10), pp. 1707–1709. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=9&issue=10&article=043>.
 42. Li, B. et al. (2020) ‘Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China’, *Clinical research in cardiology: official journal of the German Cardiac Society*, 109(5), pp. 531–538. doi: 10.1007/s00392-020-01626-9.
 43. Liu, J. et al. (2020) ‘Community Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, Shenzhen, China, 2020’, *Emerging infectious diseases*, 26(6), pp. 1320–1323. doi: 10.3201/eid2606.200239.
 44. Manohar, M. P. and Sharma, S. (2018) ‘A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists’, *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(6), pp. 716–720. doi: 10.4103/ijdr.IJDR_716_16.
 45. Masters, P. S. (2006) ‘The molecular biology of coronaviruses’, *Advances in virus research*, 66, pp. 193–292. doi: 10.1016/S0065-3527(06)66005-3.
 46. Mathew, M. G. et al. (2020) ‘Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival

- inflammation in primary molars: Randomized controlled trial', *Clinical oral investigations*, pp. 1–6. Available at: <https://link.springer.com/article/10.1007/s00784-020-03204-9>.
47. McIntosh, K. (1974) 'Coronaviruses: A Comparative Review', in *Current Topics in Microbiology and Immunology / Ergebnisse der Mikrobiologie und Immunitätsforschung*. Springer Berlin Heidelberg, pp. 85–129. doi: 10.1007/978-3-642-65775-7_3.
 48. Menon, S. et al. (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', *Colloids and surfaces. B, Biointerfaces*, 170, pp. 280–292. doi: 10.1016/j.colsurfb.2018.06.006.
 49. Nandakumar, M. and Nasim, I. (2018) 'Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis', *Journal of conservative dentistry: JCD*, 21(5), pp. 516–520. doi: 10.4103/JCD.JCD_110_18.
 50. Nandhini, J. S. T., Babu, K. Y. and Mohanraj, K. G. (2018) 'Size, shape, prominence and localization of gerdy's tubercle in dry human tibial bones', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3604. doi: 10.5958/0974-360x.2018.00663.7.
 51. Ong, S. W. X. et al. (2020) 'Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient', *JAMA: the journal of the American Medical Association*. doi: 10.1001/jama.2020.3227.
 52. Pc, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research*. Available at: <https://europepmc.org/article/med/29624863>.
 53. Prashaanthi, N. and Brundha, M. P. (2018) 'A Comparative Study between Popplet Notes and Conventional Notes for Learning Pathology', *Research Journal of Pharmacy and Technology*, 11(1), pp. 175–178. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=11&issue=1&article=032>.
 54. Preethikaa, S. and Brundha, M. P. (2018) 'Awareness of diabetes mellitus among general population', *Research Journal of Pharmacy and Technology*, 11(5), pp. 1825–1829. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=11&issue=5&article=024>.
 55. Rajakeerthi and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Üniversitesi Diş Hekimliği Fakültesi dergisi*, 22(2), pp. 249–256. doi: 10.7126/cumudj.525182.
 56. Rajendran, R. et al. (2019) 'Comparative evaluation of remineralizing potential of a paste containing bioactive glass and a topical cream containing casein phosphopeptide-amorphous calcium phosphate: An in vitro study', *Pesquisa brasileira em odontopediatria e clinica integrada*, 19(1), pp. 1–10. doi: 10.4034/pboci.2019.191.61.
 57. Ramadurai, N. et al. (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial', *Clinical oral investigations*, 23(9), pp. 3543–3550. doi: 10.1007/s00784-018-2775-5.
 58. Ramesh, A. et al. (2018) 'Comparative estimation of sulfiredoxin levels between chronic

- periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248. doi: 10.1002/JPER.17-0445.
59. Ravichandran, H. and Brundha, M. P. (2016) 'Awareness about personal protective equipments in hospital workers (sweepers and cleaners)', *International Journal of Pharmaceutical Sciences Review and Research*, 40(1), pp. 28–29.
60. Ravintha, K. and Jayalakshmi (2018) 'Recent advancements in laminates and veneers in dentistry', *Journal of advanced pharmaceutical technology & research*, 11(2), p. 785. doi: 10.5958/0974-360x.2018.00148.8.
61. R, H. et al. (2020) 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 306–312. doi: 10.1016/j.joooo.2020.06.021.
62. Saif, F. (no date) 'COVID-19 Pandemic in Pakistan: Stages and Recommendations'. doi: 10.1101/2020.05.11.20098004.
63. Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286. doi: 10.1111/ipd.12662.
64. Sekar, D. et al. (2019) 'Methylation-dependent circulating microRNA 510 in preeclampsia patients', *Hypertension research: official journal of the Japanese Society of Hypertension*, 42(10), pp. 1647–1648. doi: 10.1038/s41440-019-0269-8.
65. Seppan, P. et al. (2018) 'Therapeutic potential of Mucuna pruriens (Linn.) on ageing induced damage in dorsal nerve of the penis and its implication on erectile function: an experimental study using albino rats', *The aging male: the official journal of the International Society for the Study of the Aging Male*, pp. 1–14. doi: 10.1080/13685538.2018.1439005.
66. Sharma, P. et al. (2019) 'Emerging trends in the novel drug delivery approaches for the treatment of lung cancer', *Chemico-biological interactions*, 309, p. 108720. doi: 10.1016/j.cbi.2019.06.033.
67. Shen, K. et al. (2020) 'Diagnosis, treatment, and prevention of 2019 novel coronavirus infection in children: experts' consensus statement', *World journal of pediatrics: WJP*. doi: 10.1007/s12519-020-00343-7.
68. Shenoy, P. B. and Brundha, M. P. (2016) 'Awareness of polycystic ovarian disease among females of age group 18-30 years', *Research journal of pharmaceutical, biological and chemical sciences*, 8(8), p. 813. Available at: <http://search.proquest.com/openview/a8a09e7b2e9d2f967bf3fee479c7018a/1?pq-origsite=gscholar&cbl=54977>.
69. Shreya, S. and Brundha, M. P. (2017) 'Alteration of Haemoglobin Value in Relation to Age, Sex and Dental Diseases-A Retrospective Correlation Study', *Research Journal of Pharmacy and Technology*, 10(5), pp. 1363–1366. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=10&issue=5&article=016>.
70. Siddique, R. et al. (2019) 'Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi', *Journal of conservative dentistry: JCD*, 22(1), pp. 40–47. doi: 10.4103/JCD.JCD_284_18.

71. Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306. doi: 10.1111/jop.12835.
72. Su, S. *et al.* (2016) 'Epidemiology, Genetic Recombination, and Pathogenesis of Coronaviruses', *Trends in microbiology*, 24(6), pp. 490–502. doi: 10.1016/j.tim.2016.03.003.
73. Teja, K. V., Ramesh, S. and Priya, V. (2018) 'Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study', *Journal of conservative dentistry: JCD*, 21(6), pp. 592–596. doi: 10.4103/JCD.JCD_154_18.
74. Timothy, C. N., Samyuktha, P. S. and Brundha, M. P. (2019) 'Dental pulp Stem Cells in Regenerative Medicine--A Literature Review', *Research Journal of Pharmacy and Technology*, 12(8), pp. 4052–4056. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=8&article=088>.
75. Vijayashree Priyadharsini, J. (2019) 'In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens', *Journal of periodontology*, 90(12), pp. 1441–1448. doi: 10.1002/JPER.18-0673.
76. Vijayashree Priyadharsini, J., Smiline Girija, A. S. and Paramasivam, A. (2018) 'In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species', *Archives of oral biology*, 94, pp. 93–98. doi: 10.1016/j.anchoralbio.2018.07.001.
77. Wang, D. *et al.* (2020) 'Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China', *JAMA: the journal of the American Medical Association*, 323(11), pp. 1061–1069. doi: 10.1001/jama.2020.1585.
78. Woo, P. C. Y. *et al.* (2005) 'Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia', *Journal of virology*, 79(2), pp. 884–895. doi: 10.1128/JVI.79.2.884-895.2005.
79. Zhang, Y. *et al.* (2020) 'Isolation of 2019-nCoV from a stool specimen of a laboratory-confirmed case of the coronavirus disease 2019 (COVID-19)', *China CDC Weekly*, 2(8), pp. 123–124. Available at: <http://weekly.chinacdc.cn/cn/article/doi/10.46234/ccdw2020.033>.
80. Zhu, N. *et al.* (2020) 'A Novel Coronavirus from Patients with Pneumonia in China, 2019', *The New England journal of medicine*, 382(8), pp. 727–733. doi: 10.1056/NEJMoa2001017.