

## **Prevalence of Gingival Hyperpigmentation among Older Individuals More Than 40 Years of Age**

**Running Title :** Prevalence of gingival hyperpigmentation among older individuals.

**Type of Study:** A retrospective study.

### **T.Santhosh**

Saveetha Dental College and hospitals,  
Saveetha Institute of Medical and Technical Sciences(SIMATS),  
Saveetha university, Chennai - 600 077  
Tamilnadu, India.  
151301093.sdc@savaatha.com.

### **Dr. Kaarthikeyan .G**

Professor, Department of periodontics,  
Saveetha Dental College and hospitals,  
Saveetha Institute of Medical and Technical Sciences(SIMATS),  
Saveetha university, Chennai - 600 077  
Tamilnadu, India.  
kaarthikeyan@saveetha.com

### **Dr. Sankari malaiappan**

Professor, Department of periodontics,  
Saveetha Dental College and hospitals,  
Saveetha Institute of Medical and Technical Sciences(SIMATS),  
Saveetha university, Chennai - 600 077  
Tamilnadu, India.  
sankari@saveetha.com.

### **Corresponding author:**

### **Dr. Kaarthikeyan .G**

Professor, Department of periodontics,  
Saveetha Dental College and hospitals,  
Saveetha Institute of Medical and Technical Sciences(SIMATS),  
Saveetha university, Chennai - 600 077  
Tamilnadu, India  
kaarthikeyan@saveetha.com

### **ABSTRACT:**

Hyperpigmentation of the gingiva is caused by excessive melanin deposition by the melanocytes mainly located in the basal and suprabasal cell layers of the epithelium. It can also be caused by some systemic disorders. Sometimes pigmentations are due to the skin tone of the particular

individual. The prevalence of gingival hyperpigmentation among older individuals still remains an area of study where it should be fully investigated. The main purpose of this study is to evaluate the prevalence of gingival hyperpigmentation among individuals more than 40 years of age. Data retrieved from saveetha dental college. Diagnostic photographs both intra oral frontal and intra oral lateral photographs of 100 patients were retrieved and assessed with the help of SPSS software. The findings of this study shows that there is no age and gender correlation with pigmentation. Thus our study concludes that there is moderate to severe gingival pigmentation among older individuals and it has become one of the esthetic concerns among people. With the emergence of depigmentation treatment as one of the important array of periodontal therapy.

### **KEYWORDS:**

Esthetics; Gingival Hyperpigmentation; Melanin pigmentation.

### **INTRODUCTION:**

A smile expresses a feeling of joy, success, sensuality, affection and courtesy, and can express one's self confidence and kindness. Harmony of the smile not only depends on shape, colour and position of the teeth, but also influenced by gingival tissues. Health and appearance of gingival tissues plays a major role in an esthetic smile [(Priyanka *et al.*, 2017)]. The gingiva is the most commonly pigmented tissue of the oral cavity. Esthetics in dentistry means not only to get whiter teeth, but to create a Pink Gingiva, is always a daunting task for the treating Periodontist. Normal colour of gingiva is coral pink [(Hedin and Axéll, 1991; Hedin, Pindborg and Axéll, 1993)]. Esthetics of the intra oral soft tissue has become a significant aspect of dentistry and clinicians are faced with achieving acceptable and good gingival esthetics. On the increasing demand of esthetics, numerous procedures have been performed such as gingivectomy, gingivoplasty, connective tissue grafting, and guided tissue regeneration to improve gingival esthetics [(Tamizí and Taheri, 1996)]. Although the color of the gingiva plays an important role in overall esthetics, the principles and the techniques of the management of the problems associated with gingival melanin pigmentation (GMP) are still not established [(Ramesh *et al.*, 2019)]. Gingival hyperpigmentation is a condition in which part or whole of the gingiva is darkly pigmented in contrast to the pink or coral pink pigmentation [(Varghese *et al.*, 2015)]. It is caused by excessive deposition of melanin in the basal and suprabasal cell layers of the epithelium [(Ramesh, Sheeja S. Varghese, *et al.*, 2016)]. These melanocytes are dendritic unattached to the surrounding epithelial cells. These melanocytes when becomes active convert tyrosine to melanin protein which is then transferred to the basal and prickle cell layers [(Ramesh, Sheeja Saji Varghese, *et al.*, 2016)].

Hyperpigmentation can be caused due to a variety of local and systemic factors. Some of the Systemic conditions are endocrine disturbance, Albright's syndrome, malignant melanoma, antimalarial therapy, Peutz-Jeghers syndrome, trauma, hemochromatosis, chronic pulmonary disease [(Ramamurthy and Mg, 2018)]. Mucosal diseases in particular lichen planus can cause mucosal pigmentation which is clinically seen as multiple brown-black-pigmented areas adjacent to reticular or erosive lesions of lichen planus [(Kavarthapu and Thamaraiselvan, 2018)]. Although gingival melanin pigmentation does not represent a pathological problem, patients with a gummy smile or excessive gingival display usually complain of a "black gum" and request cosmetic therapy [(Khalid *et al.*, 2016)].

Skin tone is also a known cause of oral melanin pigmentation. High levels of oral melanin pigmentation are seen normally in individuals of African, East Asian, or Hispanic ethnicity [(Khalid *et al.*, 2017)]. In general, individuals with fair skin will not demonstrate overt tissue pigmentation, although comparable numbers of melanocytes are present within their gingival epithelium [(Schroeder, 1969)]. Smoking has been found to be associated with the provocation of the melanocytes to produce melanin [(Ravi *et al.*, 2017)]. This happens upon stimulation by polycyclic amines like nicotine and benzopyrenes that happen to be present in the tobacco, which are known to penetrate into the oral mucosa and bind to melanin [(Hedin and Axéll, 1991)]. The term “smoker’s melanosis” has been used to describe a benign focal melanin pigmentation usually present in the attached gingiva among tobacco smokers, which was described in previous literature [(Mootha *et al.*, 2016)]. This condition is not associated with any form of genetic factors, medical conditions, or the use of therapeutic medications [(Panda *et al.*, 2014)].

Classification of gingival pigmentation according to Dummett et al ranges from no pigmentation through to mild and moderate pigmentation to severe form of gingival hyperpigmentation [(Avinash, Malaippan and Dooraiswamy, 2017)]. Clinically, the hyperpigmentation can appear as a bilateral, well demarcated, ribbon-like, or dark brown band. Mucosal colour can range from light to dark brown [(Dummett and Barends, 1971)]. The attached gingiva is the most common intraoral site for melanin pigmentation and usually the marginal gingiva seems to be not affected. The buccal mucosa, hard palate, lips and tongue may also be affected [(Çiçek and Ertaş, 2003)]. The hyperpigmentation has been noted to be more in the incisor regions and decreases posteriorly. There are several management modalities for gingival pigmentation unknown to the public and dental practitioners and they comprises of gingivectomy with free gingival autografting, surgical depigmentation, electrosurgery, cryotherapy, and with the help of some lasers Nd:YAG lasers, diode laser, argon laser, CO2 laser [(Moravej-Salehi, Moravej-Salehi and Hajifattahi, 2015)].

Our team has rich experience in research and we have collaborated with numerous authors over various topics in the past decade (Ariga *et al.*, 2018; Basha, Ganapathy and Venugopalan, 2018; Hannah *et al.*, 2018; Hussainy *et al.*, 2018; Jeevanandan and Govindaraju, 2018; Kannan and Venugopalan, 2018; Kumar and Antony, 2018; Manohar and Sharma, 2018; Menon *et al.*, 2018; Nandakumar and Nasim, 2018; Nandhini, Babu and Mohanraj, 2018; Ravinthar and Jayalakshmi, 2018; Seppan *et al.*, 2018; Teja, Ramesh and Priya, 2018; Duraisamy *et al.*, 2019; Gheena and Ezhilarasan, 2019; Hema Shree *et al.*, 2019; Rajakeerthi and Ms, 2019; Rajendran *et al.*, 2019; Sekar *et al.*, 2019; Sharma *et al.*, 2019; Siddique *et al.*, 2019; Janani, Palanivelu and Sandhya, 2020; Johnson *et al.*, 2020; Jose, Ajitha and Subbaiyan, 2020).

There is insufficient knowledge on prevalence of gingival hyperpigmentation in tamilnadu. Particularly prevalence of gingival hyperpigmentation among older individuals still remains to be fully investigated. The findings from this study can help dentists to gain knowledge about gingival pigmentation.

## **MATERIALS AND METHOD:**

The case records of 86000 patients visiting Saveetha Dental College were analysed and a total of 100 patients with gingival pigmentation were analysed. All the data of 100 patients who

visited saveetha dental college were retrieved retrospectively. Ethical clearance: Ethical clearance was obtained from the research ethical board of saveetha dental college prior to the study. Sample size was estimated to be 100 patients who visited saveetha dental college between June 2019 to April 2020 were selected for review. The case sheets were analysed for the following variables of age, gender, diagnostic photographs were analysed. In case of doubts or discordance of Data, the patients were contacted over the phone or asked to report back to the College to confirm the findings.

**Inclusion criteria:**

- Patients above 40 years of age
- Proper diagnostic photographs with proper lighting

**Exclusion criteria:**

- Patient with other oral lesions.
- Patients below the age of 40 years.

All the statistics and analysis were done using SPSS software (version 2019). All the descriptive analysis such as mean standard deviation and percentages were used to present the number of male and female subjects and demographic variables. Chi square test was used to establish correlation between categorical variables.  $P (<0.05)$  was set to be statistically significant.

The internal validity of the study was established as the data was collected from a verifiable and standardised database. The external validity is established as the data is from a clinical setup which is duplicatable.

**RESULTS:**

Among all the 100 subjects. More number of subjects were found to be in the range between 41 to 50 years of age with 48% (Figure 1). No of males ( $n=57$ ) and females ( $n=43$ ) (Figure 2). Qualitative analysis of pigmentation: 100 subjects were evaluated and indexed using dummett et al classification and found patients above 40 year of age have a high prevalence of gingival hyperpigmentation. Mild, moderate and severe accounts ( $n=74$ ) of total 100 subjects evaluated. Only 26 subjects had no gingival pigmentation (Figure 3)(4,5,6,7). Gender predilection of pigmentation: Both male and female showed equal percentages of cases with no, mild, moderate, severe pigmentation, however male had more numbers of moderate pigmentation (19%). Pearson chi square test was done  $p=0.233 (>0.05)$  not statistically significant. No gender predominance is found in the prevalence of gingival hyperpigmentation (Figure 8). Association between age and gingival pigmentation: shows that the age group of 41-50 have more number of cases with severe pigmentation (16%). Association was assessed by chi square test;  $p=0.117 (>0.05)$ , not statistically significant.

**DISCUSSION:**

Melanin pigmentation often occurs in the gingiva as a result of an abnormal or increased deposition of melanin [(Ramesh, Ravi and Kaarthikeyan, 2017)]. Brown or dark pigmentation and discoloration of gingival tissue, whether of a physiologic or pathologic nature, can be caused by a variety of local and systemic factors. This type of pigmentation is symmetric and it is found to be persistent, and it does not alter normal structures of gingiva. This pigmentation may be seen across all races and at any age, and has no gender predilection [(Thamaraiselvan *et al.*, 2015)].

Dummett proposed that the degree of pigmentation is directly related to the mechanical, chemical, physiologic stimulators of melanocytes. Fair skinned and dark skinned people have the same levels of melanocytes but, the difference depends upon the activity of melanocytes [(Haresaku *et al.*, 2007)]. It has been reported that gingiva is the most commonly pigmented site in the oral cavity and children are less affected than adults. Pigmentation is caused by physiologic pigmentation and it causes an esthetic concern for the adults leading to formation of “Black Gingiva”. Pigmentation can be seen due to drugs, smoking, heavy metals, bronze diabetes, HIV infection and other diseases. In this study we analysed the gingival pigmentation in 100 patients from 40 to 90 years of age. We found (n=51) half of the study subjects belongs to the age group of 41-50 years of age. (n=27) belongs to the age group of 51-60 years of age. And fewer number of subjects from the rest of the age groups. Gingival pigmentation is seen as a genetic trait in some populations [(Rosa *et al.*, 2007)]. The prevalence of melanin pigmentation varies between 0% to 89%. Secondary influences can be smoking [(Dummett and Barens, 1971)].

Among the same ethnic people or race the pigmentation depends on megaloblastic activity [(Prasad, Agrawal and Reddy, 2010)]. Melanin, eumelanin, pheomelanin, and neuromelanin are the common types. In general people who lived for longer periods have more pigmentation [(Ramesh, Sheeja S. Varghese, *et al.*, 2016)]. The distribution of gingival pigmentation is unique in south indians. In this study we observed patients above 40 years of age have moderate to severe pigmentation [(Prasad *et al.*, 2005)]. A Particular age range of (41-50) years shows severe pigmentation which contradicts the previous studies. This study also shows there is no particular difference between male and female gender which is similar to the previous studies [(Araki *et al.*, 1983)]. In this present study age is correlated with pigmentation generally and there was no specific correlation seen. These findings could be attributed that there is no variation within a specific ethnic group. Small sample size and no direct patient evaluation were found to be the limitations of this study.

Our institution is passionate about high quality evidence based research and has excelled in various fields ( (Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; Vijayashree Priyadharsini, Smiline Girija and Paramasivam, 2018; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Ramadurai *et al.*, 2019; Sridharan *et al.*, 2019; Vijayashree Priyadharsini, 2019; Chandrasekar *et al.*, 2020; Mathew *et al.*, 2020; R *et al.*, 2020; Samuel, 2021)

## CONCLUSION:

Gingival pigmentation has become one of the concerns among people. With the emergence of depigmentation treatment as one of the important array of periodontal therapy. The findings of this study conclude that there is no age and gender correlation with pigmentation. More studies with larger patients samples needed to be done in future including skin tone, smoking habits in this specific ethnic group.

## AUTHOR CONTRIBUTIONS:

**Santosh** had contributed to the design of the study, data collection, analysis of data, results tabulation, manuscript preparation.

**Kaarthikeyan** had contributed to the design of the study, analysis of data, results, manuscript preparation.

**Sankari** had contributed to the design of the study, manuscript preparation, proofreading of the manuscript.

#### **CONFLICT OF INTEREST:**

This research project is self funded and is not sponsored or aided by any third party. There is no conflict of interest.

#### **REFERENCES:**

- [1]. Araki, S. *et al.* (1983) 'Dose-response relationship between tobacco consumption and melanin pigmentation in the attached gingiva', *Archives of environmental health*, 38(6), pp. 375–378. doi: 10.1080/00039896.1983.10545823.
- [2]. Ariga, P. *et al.* (2018) 'Determination of correlation of width of Maxillary Anterior Teeth using Extraoral and Intraoral Factors in Indian Population: A systematic review', *World journal of dentistry*, 9(1), pp. 68–75. doi: 10.5005/jp-journals-10015-1509.
- [3]. Avinash, K., Malaippan, S. and Dooraiswamy, J. N. (2017) 'Methods of Isolation and Characterization of Stem Cells from Different Regions of Oral Cavity Using Markers: A Systematic Review', *International journal of stem cells*, 10(1), pp. 12–20. doi: 10.15283/ijsc17010.
- [4]. Basha, F. Y. S., Ganapathy, D. and Venugopalan, S. (2018) 'Oral hygiene status among pregnant women', *Journal of advanced pharmaceutical technology & research*, 11(7), p. 3099. doi: 10.5958/0974-360x.2018.00569.3.
- [5]. Chandrasekar, R. *et al.* (2020) 'Development and validation of a formula for objective assessment of cervical vertebral bone age', *Progress in orthodontics*, 21(1), p. 38. doi: 10.1186/s40510-020-00338-0.
- [6]. Çiçek, Y. and Ertaş, U. (2003) 'The normal and pathological pigmentation of oral mucous membrane: a review', *The journal of contemporary dental practice*, 4(3), pp. 76–86. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12937598>.
- [7]. Dummett, C. O. and Barens, G. (1971) 'Oromucosal pigmentation: an updated literary review', *Journal of periodontology*, 42(11), pp. 726–736. doi: 10.1902/jop.1971.42.11.726.
- [8]. Duraisamy, R. *et al.* (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295. doi: 10.1097/ID.0000000000000885.
- [9]. Ezhilarasan, D., Apoorva, V. S. and Ashok Vardhan, N. (2019) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121. doi: 10.1111/jop.12806.
- [10]. Gheena, S. and Ezhilarasan, D. (2019) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & experimental toxicology*, 38(6), pp. 694–702. doi: 10.1177/0960327119839173.
- [11]. Hannah, R. *et al.* (2018) 'Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens', *Journal of advanced pharmaceutical technology & research*, 11(3), p. 1012. doi: 10.5958/0974-360x.2018.00189.0.
- [12]. Haresaku, S. *et al.* (2007) 'Association of lip pigmentation with smoking and gingival melanin pigmentation', *Oral diseases*, 13(1), pp. 71–76. doi: 10.1111/j.1601-0825.2006.01249.x.

- [13]. Hedin, C. A. and Axéll, T. (1991) 'Oral melanin pigmentation in 467 Thai and Malaysian people with special emphasis on smoker's melanosis', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 20(1), pp. 8–12. doi: 10.1111/j.1600-0714.1991.tb00879.x.
- [14]. Hedin, C. A., Pindborg, J. J. and Axéll, T. (1993) 'Disappearance of smoker's melanosis after reducing smoking', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 22(5), pp. 228–230. doi: 10.1111/j.1600-0714.1993.tb01061.x.
- [15]. Hema Shree, K. *et al.* (2019) 'Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma - a Systematic Review with Meta Analysis', *Pathology oncology research: POR*, 25(2), pp. 447–453. doi: 10.1007/s12253-019-00588-2.
- [16]. Hussainy, S. N. *et al.* (2018) 'Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up', *Journal of conservative dentistry: JCD*, 21(5), pp. 510–515. doi: 10.4103/JCD.JCD\_51\_18.
- [17]. Janani, K., Palanivelu, A. and Sandhya, R. (2020) 'Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality: an in vivo study', *Brazilian dental science*, 23(1). doi: 10.14295/bds.2020.v23i1.1805.
- [18]. Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 19(4), pp. 273–278. doi: 10.1007/s40368-018-0356-6.
- [19]. Johnson, J. *et al.* (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH', *Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362. doi: 10.1038/s41440-019-0369-5.
- [20]. Jose, J., Ajitha and Subbaiyan, H. (2020) 'Different treatment modalities followed by dental practitioners for Ellis class 2 fracture – A questionnaire-based survey', *The open dentistry journal*, 14(1), pp. 59–65. doi: 10.2174/1874210602014010059.
- [21]. Kannan, A. and Venugopalan, S. (2018) 'A systematic review on the effect of use of impregnated retraction cords on gingiva', *Journal of advanced pharmaceutical technology & research*, 11(5), p. 2121. doi: 10.5958/0974-360x.2018.00393.1.
- [22]. Kavarthapu, A. and Thamaraiselvan, M. (2018) 'Assessing the variation in course and position of inferior alveolar nerve among south Indian population: A cone beam computed tomographic study', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(4), pp. 405–409. doi: 10.4103/ijdr.IJDR\_418\_17.
- [23]. Khalid, W. *et al.* (2016) 'Role of endothelin-1 in periodontal diseases: A structured review', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 27(3), pp. 323–333. doi: 10.4103/0970-9290.186247.
- [24]. Khalid, W. *et al.* (2017) 'Comparison of Serum Levels of Endothelin-1 in Chronic Periodontitis Patients Before and After Treatment', *Journal of clinical and diagnostic research: JCDR*, 11(4), pp. ZC78–ZC81. doi: 10.7860/JCDR/2017/24518.9698.
- [25]. Kumar, D. and Antony, S. D. P. (2018) 'Calcified canal and negotiation-A review', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3727. doi: 10.5958/0974-360x.2018.00683.2.

- [26]. Manohar, M. P. and Sharma, S. (2018) 'A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(6), pp. 716–720. doi: 10.4103/ijdr.IJDR\_716\_16.
- [27]. Mathew, M. G. *et al.* (2020) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: Randomized controlled trial', *Clinical oral investigations*, pp. 1–6. Available at: <https://link.springer.com/article/10.1007/s00784-020-03204-9>.
- [28]. Menon, S. *et al.* (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', *Colloids and surfaces. B, Biointerfaces*, 170, pp. 280–292. doi: 10.1016/j.colsurfb.2018.06.006.
- [29]. Mootha, A. *et al.* (2016) 'The Effect of Periodontitis on Expression of Interleukin-21: A Systematic Review', *International journal of inflammation*, 2016. doi: 10.1155/2016/3507503.
- [30]. Moravej-Salehi, E., Moravej-Salehi, E. and Hajifattahi, F. (2015) 'Relationship of Gingival Pigmentation with Passive Smoking in Women', *Tanaffos*, 14(2), pp. 107–114. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26528364>.
- [31]. Nandakumar, M. and Nasim, I. (2018) 'Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis', *Journal of conservative dentistry: JCD*, 21(5), pp. 516–520. doi: 10.4103/JCD.JCD\_110\_18.
- [32]. Nandhini, J. S. T., Babu, K. Y. and Mohanraj, K. G. (2018) 'Size, shape, prominence and localization of gerdy's tubercle in dry human tibial bones', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3604. doi: 10.5958/0974-360x.2018.00663.7.
- [33]. Panda, S. *et al.* (2014) 'Platelet rich fibrin and xenograft in treatment of intrabony defect', *Contemporary clinical dentistry*, 5(4), pp. 550–554. doi: 10.4103/0976-237X.142830.
- [34]. Pc, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research*. Available at: <https://europepmc.org/article/med/29624863>.
- [35]. Prasad, D. *et al.* (2005) 'Treatment of gingival pigmentation : A case series', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 16(4), p. 171. doi: 10.4103/0970-9290.29901.
- [36]. Prasad, S. S., Agrawal, N. and Reddy, N. R. (2010) 'Gingival depigmentation: A case report', *People's J Sci Res*, 3(1), pp. 27–29. Available at: <https://www.pjsr.org/abstract-PDF/Dr.%20Neeraj%20Agrawal.pdf>.
- [37]. Priyanka, S. *et al.* (2017) 'Detection of cytomegalovirus, Epstein-Barr virus, and Torque Teno virus in subgingival and atheromatous plaques of cardiac patients with chronic periodontitis', *Journal of Indian Society of Periodontology*, 21(6), pp. 456–460. doi: 10.4103/jisp.jisp\_205\_17.
- [38]. Rajakeerthi and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Üniversitesi Diş Hekimliği Fakültesi dergisi*, 22(2), pp. 249–256. doi: 10.7126/cumudj.525182.
- [39]. Haque, M. Handwashing in averting infectious diseases: Relevance to COVID-19, (2020) *Journal of Population Therapeutics and Clinical Pharmacology*, 27 (Special Issues 1), pp. e37-e52.
- [40]. Ramadurai, N. *et al.* (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial', *Clinical oral investigations*, 23(9), pp. 3543–3550. doi: 10.1007/s00784-018-2775-5.
- [41]. Ramamurthy, J. and Mg, V. (2018) 'COMPARISON OF EFFECT OF HIORA MOUTHWASH VERSUS CHLORHEXIDINE MOUTHWASH IN GINGIVITIS PATIENTS: A CLINICAL



- TRIAL', *Asian J Pharm Clin Res*, 11(7), pp. 84–88. Available at: <https://pdfs.semanticscholar.org/1c22/6e98fc99e9fb99bc749ae5d553024fa93052.pdf>.
- [42]. Ramesh, A., Varghese, S. S., *et al.* (2016) 'Chronic obstructive pulmonary disease and periodontitis--unwinding their linking mechanisms', *Journal of oral biosciences / JAOB, Japanese Association for Oral Biology*, 58(1), pp. 23–26. Available at: <https://www.sciencedirect.com/science/article/pii/S1349007915001103>.
- [43]. Ramesh, A., Varghese, S. S., *et al.* (2016) 'Herbs as an antioxidant arsenal for periodontal diseases', *Journal of intercultural ethnopharmacology*, 5(1), pp. 92–96. doi: 10.5455/jice.20160122065556.
- [44]. Ramesh, A. *et al.* (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248. doi: 10.1002/JPER.17-0445.
- [45]. Ramesh, A. *et al.* (2019) 'Esthetic lip repositioning: A cosmetic approach for correction of gummy smile - A case series', *Journal of Indian Society of Periodontology*, 23(3), pp. 290–294. doi: 10.4103/jisp.jisp\_548\_18.
- [46]. Ramesh, A., Ravi, S. and Kaarthikeyan, G. (2017) 'Comprehensive rehabilitation using dental implants in generalized aggressive periodontitis', *Journal of Indian Society of Periodontology*, 21(2), pp. 160–163. doi: 10.4103/jisp.jisp\_213\_17.
- [47]. Ravinthar, K. and Jayalakshmi (2018) 'Recent advancements in laminates and veneers in dentistry', *Journal of advanced pharmaceutical technology & research*, 11(2), p. 785. doi: 10.5958/0974-360x.2018.00148.8.
- [48]. Ravi, S. *et al.* (2017) 'Additive Effect of Plasma Rich in Growth Factors With Guided Tissue Regeneration in Treatment of Intrabony Defects in Patients With Chronic Periodontitis: A Split-Mouth Randomized Controlled Clinical Trial', *Journal of periodontology*, 88(9), pp. 839–845. doi: 10.1902/jop.2017.160824.
- [49]. R, H. *et al.* (2020) 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 306–312. doi: 10.1016/j.o.oooo.2020.06.021.
- [50]. Rosa, D. S. A. *et al.* (2007) 'Esthetic treatment of gingival melanin hyperpigmentation with Er: YAG laser: Short-term clinical observations and patient follow-up', *Journal of periodontology*, 78(10), pp. 2018–2025. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1902/jop.2007.070041>.
- [51]. Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286. doi: 10.1111/ipd.12662.
- [52]. Schroeder, H. E. (1969) 'Melanin containing organelles in cells of the human gingiva I. Epithelial Melanocytes', *Journal of periodontal research*, 4(1), pp. 1–18. doi: 10.1111/j.1600-0765.1969.tb01940.x.
- [53]. Sekar, D. *et al.* (2019) 'Methylation-dependent circulating microRNA 510 in preeclampsia patients', *Hypertension research: official journal of the Japanese Society of Hypertension*, 42(10), pp. 1647–1648. doi: 10.1038/s41440-019-0269-8.
- [54]. Seppan, P. *et al.* (2018) 'Therapeutic potential of *Mucuna pruriens* (Linn.) on ageing induced damage in dorsal nerve of the penis and its implication on erectile function: an experimental study using albino rats', *The aging male: the official journal of the International Society for the Study of the Aging Male*, pp. 1–14. doi: 10.1080/13685538.2018.1439005.

- [55]. Sharma, P. *et al.* (2019) 'Emerging trends in the novel drug delivery approaches for the treatment of lung cancer', *Chemico-biological interactions*, 309, p. 108720. doi: 10.1016/j.cbi.2019.06.033.
- [56]. Siddique, R. *et al.* (2019) 'Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi', *Journal of conservative dentistry: JCD*, 22(1), pp. 40–47. doi: 10.4103/JCD.JCD\_284\_18.
- [57]. Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306. doi: 10.1111/jop.12835.
- [58]. Tamizí, M. and Taheri, M. (1996) 'Treatment of severe physiologic gingival pigmentation with free gingival autograft', *Quintessence international*, 27(8). Available at: <http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=00336572&AN=38506436&h=FvDGOpcezA5%2FW%2B0JXL254OXI9GX%2FmlyKaF8%2BqH8kpPIAFdHw8B5TChm621Ejfh8xxau8%2BHIHIK%2B8Cy6aNi9qYg%3D%3D&crl=c>.
- [59]. Teja, K. V., Ramesh, S. and Priya, V. (2018) 'Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study', *Journal of conservative dentistry: JCD*, 21(6), pp. 592–596. doi: 10.4103/JCD.JCD\_154\_18.
- [60]. Thamaraiselvan, M. *et al.* (2015) 'Comparative clinical evaluation of coronally advanced flap with or without platelet rich fibrin membrane in the treatment of isolated gingival recession', *Journal of Indian Society of Periodontology*, 19(1), pp. 66–71. doi: 10.4103/0972-124X.145790.
- [61]. Varghese, S. S. *et al.* (2015) 'Estimation of salivary tumor necrosis factor-alpha in chronic and aggressive periodontitis patients', *Contemporary clinical dentistry*, 6(Suppl 1), pp. S152–6. doi: 10.4103/0976-237X.166816.
- [62]. Vijayashree Priyadharsini, J. (2019) 'In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens', *Journal of periodontology*, 90(12), pp. 1441–1448. doi: 10.1002/JPER.18-0673.
- [63]. Vijayashree Priyadharsini, J., Smiline Girija, A. S. and Paramasivam, A. (2018) 'In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species', *Archives of oral biology*, 94, pp. 93–98. doi: 10.1016/j.archoralbio.2018.07.001.

**Tables and graphs:**




Figure 1: shows distribution of subjects among different age groups. Blue colour represents- 41-50years (49%); green represents- 51-60years (27%); sandal represents- 61-70years (16%); purple represents- 71-80years (6%); yellow represents- 81-90 (2%). Figure I inferred that the majority of the study subjects are from the age group of 41-50 years and least are from 81-90 years of age.




Figure 2: shows gender distribution of subjects. Blue colour represents- male (57%); green colour represents- female (43%). Figure II inferred that more numbers of male were included in this study than females.




Figure 3: Bar chart shows different types of pigmentation. (X axis represents- different score of pigmentation; Y axis represents- number of cases). Pink colour represents- no pigmentation; blue represents- mild pigmentation; grey represents- moderate pigmentation; purple represents- severe pigmentation. Figure II inferred that the patients above 40 years of age had moderate gingival pigmentation in more numbers, followed by no pigmentation, severe, and mild pigmentation.




Figure 4: Bar chart shows number of cases with no pigmentation in each age group. (X axis represents- age group of the subjects; Y axis represents number of cases with no pigmentation) Figure IV infrared that more number of cases with no pigmentation are from 51-60 years of age.




Figure 5: Bar chart shows number of cases with mid pigmentation in each age group. (X axis represents- age group of the subjects; Y axis represents number of cases with mild pigmentation) Figure IV infrared that more number of cases with mild pigmentation are from 41-50 years of age.




Figure 6: Bar chart shows number of cases with moderate pigmentation in each age group. (X axis represents- age group of the subjects; Y axis represents number of cases with moderate pigmentation) Figure IV infrared that more number of cases with moderate pigmentation are from 41-50 years of age. Out of 6 total subjects in the age group of 71-80years were found with moderate pigmentation.






Figure 7: Bar chart shows number of cases with severe pigmentation in each age group. (X axis represents- age group of the subjects; Y axis represents number of cases with severe pigmentation) Figure IV infrared that more number of cases with severe pigmentation are from 41-50 years of age.




Figure 8: Bar chart shows association between gender and gingival pigmentation. (X axis represents- gender of the subjects; Y axis represents percentage of cases) Both male and female showed equal percentages of cases with no, mild, moderate, severe pigmentation, however male had more numbers of moderate pigmentation (19%). Chi square test shows no significant association between gender and gingival pigmentation ,Pearson chi square test was done  $p=0.233$  ( $>0.05$ ) not statistically significant.




Figure 9: Bar graph represents association between age and gingival pigmentation. (X axis represents- age of the subjects; Y axis represents percentage of cases) subjects from the age group of 41-50 have more number of cases with severe pigmentation (16%, purple colour), subjects from the age group of 51-60 had more number of cases with no pigmentation (11%, blue colour). Chi square test shows no significant association between gender and gingival pigmentation ,Pearson chi square test was done  $p=0.117 (>0.05)$  not statistically significant.