

The Effectiveness of the Integration among *Beauveria bassiana* Vuill., *Isaria fumosorosea* (Wize) and Insecticides Against *Bemisia tabaci* (Genn.)

¹ Asst. Lect. Ridhab Faleh Hassan , ² Prof. Dr Bassim Sh. Hamad,
³ Prof. Dr. Mohammed J. Hanawi.

¹University of Wasit, College of Science

² Ministry of Science and Technology, Directorate of Agric. Res, IPM Center, Iraq,

³ University of Wasit, College of Science, Iraq

83ridab83@gmail.com, bassim.shehab@yahoo.com, hanawimohammed@yahoo.com

Corresponding Author: 83ridab83@gmail.com

Abstract

This research was carried out to determine the compatibility of different components of integrated pest management (the entomopathogenic fungi: *Isaria fumosorosea*, *Beauveria bassiana* Bbk, the pesticide Imidacloprid and Mineral oil) and their efficiency to control the white fly *Bemisia tabaci* on tomato. The results showed a weak effect of the two pesticides at a field dose and a half field dose on the vegetative growth and spores production of *B. bassiana* Bbk and *I. fumosorosea*, this reflects the compatibility between the two pesticides and these fungi. Using of their combinations against whitefly nymphs showed that the Imidacloprid + *I. fumosorosea* gave the highest mortality rate after three days of treatment (79.6%), followed by the Mineral oil with *B. bassiana* Bbk (78.6%). After seven days the superiority of the Imidacloprid + *I. fumosorosea* continued (91.7 %) without significant difference from other combinations, but with significant difference from the agents used alone.

Key words: *Isaria fumosorosea*, *Beauveria bassiana*, Imidacloprid, compatibility, *Bemisia tabaci*.

Introduction

Bemisia tabaci (Genn.) (Hemiptera: Aleyrodidae) is one of the most damaging and common insect pests of greenhouse and field crops around the world. Due to direct damage and the spread of many plant viruses, it poses a significant danger to crop production ⁽¹⁾. Chemical insecticides are used mainly to combat this insect, but this process has run into issues due to the selection of resistant individuals. Furthermore, their adverse side effects on non-target species, as well as their environmental risks, have caused the use of alternate monitoring measures. As a result, the search for more environmentally friendly and long-lasting alternate management approaches opens up new ways to study microbial agents with biological control capacity. As a consequence, entomopathogenic fungi refer to a category of microorganisms

<http://annalsofrscb.ro>

that spontaneously infect and thereby control whitefly populations. Agents of biological regulation Scientists, society, and private businesses are all interested in microbial pathogens of arthropods because they provide an environmentally safe alternative to conventional pesticides for use in advanced pest management programs (IPM) ⁽²⁾. *Beauveria bassiana* (Bals-Criv.) Vuill., *Isaria fumosorosea* (Wize) (formerly *Paecilomyces fumosoroseus*), *Lecanicillium* spp. (formerly *Verticillium* spp.), and *Aschersonia* spp. are among the most promising fungal candidates for whitefly biological control. ⁽³⁾. The aim of this analysis was to see how successful *Isaria fumosorosea*, *Beauveria bassiana*, Imidacloprid, and Mineral oil were at controlling the whitefly *Bemisia tabaci*.

Methodology

The whitefly *Bemisia tabaci*, which used in the study, was obtained by infecting tomato plants (Super queen variety) by using infected tomato's leaves collected from the fields. They were planted in 10 cm diameter pots with a mixture of soil and peat moss (1:1) and were maintained under a green house condition. Furthermore, two local isolates (*Beauveria bassiana* (Bbk) and *Isaria fumosorosea*.) are used in this study, Obtained from the Ministry of Science and Technology', Directorate of Agricultural Research. The following chemical pesticides were used:

- 1- Bright 25% (wp), the active ingredient is Imidacloprid, the field dose rate is 1 g / liter of water
- 2- Mineral oil Nautilus 70% EW , the field rate is 20 ml / liter of water.

Pathogenic activity of *B. bassiana* Bbk and *I. fumosorosea* was measured at concentrations of 10^5 , 10^7 and 10^9 spores / ml + 0.01% Tween 20 on whitefly nymphs by spraying three plants for each concentration (treatment), the control treatment was sprayed with water +0.01% Tween 20. The chemical pesticides were used with the concentration of field and half field doses. The mortality rate of the nymphs was counted after 3, 5 and 7 days of treatment with fungi and three days for chemical pesticides that corrected using Abbott's formula ⁽⁴⁾.

The tests for fungal vegetative growth, and conidia production were done. Insecticides used in this study were imidacloprid and Nautilus mineral oil, with two doses, field and half field doses. Vegetative growth was measured by poisoned food technique ⁽⁵⁾. The formula suggested by (6) was used to identify chemical products based on mycelia growth (VG) and sporulation (SP) in addition to the regulation in in vitro experiments with entomopathogenic fungi (100 percent). The index is less than 30 in $T = 20 (VG) + 80 (SP) / 100$: Toxic.

An experiment of integration between fungi and chemical pesticides was carried out on tomato plants (Super queen variety) grown in pots, the pesticide was sprayed and the next day the spore suspension was sprayed with fungi. The experiment was designed according to

randomized complete block design with three blocks, each with one plant for each treatment under greenhouse conditions. The treatments included in the study are shown in Table (1).

Table (1): Treatments

Treatments	Dose rate
Imidocloprid (Im)	half field dose
Im + <i>Isaria</i>	half field dose +10^7
Im + <i>Bbk</i>	half field dose +10^7
mineral oil	half field dose
Mineral oil +<i>Isaria</i>	half field dose +10^7
Mineral Oil + <i>Bbk</i>	half field dose +10^7
<i>Isaria</i>	10^7
<i>Bbk</i>	10^7


Results and Discussion

The tests were planned using a randomized full block method (RCBD), and the data was evaluated using the SPSS software and the Duncan multiple range test (DMRT) to compare the means at the 0.05 likelihood stage. Standard probit analysis was used to obtain LC50 and LT50 values.

The impact of *B. bassiana* *Bbk* and *I. fumosorosea* fungal suspensions on whitefly nymphs was related to isolate and concentrations. The highest mortality rate was achieved after seven days of treatment for a concentration of 10^9 spores / ml, 65.6% of isolate *Bbk* and 69%. for isolate *I. fumosorosea*, the correlation coefficient between concentration and mortality rate was 0.93 and 0.95 for the two isolates, respectively, and by linear equation was $y = 4.6x + 29.767$ for *I. Fumosoros* and $y = 3x + 39.833$ for *B. bassiana* (Fig. 1, 2).

The pathogenicity of the fungal isolates on the basis of the LC50 were the highest ability of infection by the isolate *Bbk*, as the value of LC50 was 9.7×10^3 spores / ml after seven days of treatment, while the isolate *I. fumosorosea* recorded 2.7×10^4 spores / ml . The virulence

of isolates represented by (LT50), was inversely related to the concentration, the concentration 10^9 spores / ml given the shortest time. The isolation *Bbk* was the most virulent with the shortest time (3.23 days), followed by isolation *I. fumosorosea* (3.8 days) at the same concentration as it is clear in Figure (1) and Figure (2).



Figure(1): Linear equation of the effect of *B. bassiana*




Figure (2): Linear equation for the effect of *Isaria fumosorosea*

Table (2): LC50 and LT50 of *B. Bassiana* and *I. Fumosorosea* Against Whitefly Nymphs

Isolate	Concentrations	LT50(day)
<i>Bbk</i>	10^5	5.55
	10^7	4.0
	10^9	3.23
LC ₅₀	$10^{3.9.7 \times}$	
<i>Isf</i>	10^5	6.05
	10^7	4.33
	10^9	3.8
LC ₅₀	2.7×10^4	

The normal death of whitefly species is highly affected by entomopathogenic fungi, and are distinguished by their importance as biological control agents because they can infect insects directly through the body wall ⁽²⁾. Screening of the entomopathogenic fungi to select the optimum is important in an integrated pest control strategy. The most widely used species in managing *B. tabaci* are *Beauveria bassiana*, *Metarhiziumanisopliae*, *Isariafumosorosea*, *Ashersonia* spp., and *Verticilliumlecanii*. ⁽³⁾several studies confirmed the effectiveness of *B. bassiana* in controlling *B. tabaci* whiteflies ⁽⁶⁾. Testing of twenty-five isolates of *B. bassiana* at a concentration of 10^7 spores/ml, commercial biocides of *B. bassiana* against *B. tabaci* nymph reported a mortality rate of 3-85%. The LC₅₀ of the four most virulent isolates ranged from 1.1×10^5 to 6.2×10^6 spores / ml in the second set of biological experiments on 10 isolates chosen at a concentration of 105-108 spores / ml ⁽⁷⁾.

Numerous studies confirm the importance of *Isariafumosorosea* (*Paecilomyces fumosorosea*) in controlling whiteflies. It is one of the pest's most powerful natural enemies. It can transmit *B. tabaci* epidemics in greenhouses and open fields ⁽⁸⁾.and infect a number of pests in agriculture and forest areas ⁽⁹⁾.

Under laboratory conditions, the virulence of five isolates of *B. Bassiana* and *I. fumosorosea* and four isolates of *Lecanicillium muscarium* on whiteflies on bean leaves showed that *B. bassiana* and *I. fumosorosea* were more successful, with a mortality rate of 71-86 percent within 8 days and a $Lt50$ of 3- 4 days with a concentration of 10^7 spores / ml (150 spores⁽²⁾).

Imidacloprid recorded a significant effect on nymphs of whiteflies, as the mortality rate was 91.6% at field dose, with a significant difference from the field dose of mineral oil (82.6%), and the two treatments differed significantly from the results of the half field dose of the two pesticides (75.3 and 71.3%, respectively)

Chemical control with insecticides has been the most effective way to reduce whitefly damage to crop production. Imidacloprid is one of the neoneotinoids pesticides used extensively to control pests, and it is a systemic chloronicotinyl pesticide that is used in the treatment of soil, seeds and leaves to control many pests, especially sucking insects⁽¹⁰⁾, as found by⁽¹¹⁾, Imidacloprid significantly reduced both adults and nymphs of whiteflies. Effect of Imidacloprid on whiteflies under laboratory and the greenhouse condition showed that the pesticide remains effective for 45 days⁽¹²⁾. The foliar application of Actara and Imidacloprid was substantially successful after one day of treatment on nymphs and whiteflies on eggplant, according to an assessment of the effectiveness of certain Neotinoid pesticides on nymphs and whiteflies on eggplant.

The results showed a weak effect of the two pesticides Imidacloprid and Mineral oil at a field and half field doses on *Beauveria bassiana* vegetative development and sporulation, this reflects the compatibility between the two pesticides and the fungus by 73.46 and 82.75 for Imidacloprid and 87.12 and 94.36 for the Mineral oil at Field and half doses respectively.

The reduction effect of the two pesticides on the formation of spores at the recommended field dose was 33.3% for Imidacloprid and 13.79% for mineral oil, and the reduction ratio in the half dose was 19.5 and 5.8% respectively. See Table (3):

Table (3): Pesticide Effects on Beauveria bassiana Vegetative Development and Sporulation

Treatment	Dose rate	Radius of Vegetative growth (cm)	Reduction (%)	Sporulation ($\times 10^7$)	Reduction (%)	Toxicity value	Compatibility
Imidacloprid 25%wp	Imidacloprid 25%wp	3.83	4.25	58	33.3	73.46	Compatible
	0.5 g /l	3.86	3.5	70	19.5	82.75	Compatible
Mineral oil	20ml/l	3.63	9.25	75	13.79	87.12	Compatible
	10 ml/l	3.8	5.0	82	5.8	94.36	Compatible
Control	—	4.0	—	87			

Smaller than 30: very toxic; 31-45: toxic; 46-60: moderate toxicity; greater than 60 = compatible

The two pesticides were compatible with *Isaria fumosorosea* at 66.37 and 61.37% for field and half field doses of imidacloprid respectively and by 70.26 and 76.58% for the mineral oil (Table 3), with a decrease of 29.25% and 28.25% in fungal growth and 34.48% and 40.22% for sporulation in the case of the pesticide Imidacloprid, and a decrease of 33.75% for fungal growth and 28.73 and 21.84% for sporulation in the case of mineral oil.

Table (4): Pesticide Effects on Isariafumosoro-sea vegetative Development and Sporulation

Treatment	Dose rate	Radius of Vegetative growth (cm)	Reduction (%)	Sporulation ($\times 10^7$)	Reduction (%)	Toxicity value	Compatibility
Imidacloprid 25%wp	1 g /l	2.83	29.25	52	34.48	61.37	Compatible
	0.5 g /l	2.87	28.25	57	40.22	66.37	Compatible
Mineral oil	20ml/l	2.65	33.75	62	21.84	76.58	Compatible
	10 ml/l	2.65	33.75	68	28.73	70.26	Compatible
Control			—	87			

Smaller than 30: very toxic; 31-45: toxic; 46-60: moderate toxicity; greater than 60 = compatible

The compatibility between entomopathogenic fungi and the pesticides is essential in integrated pest management programs ⁽¹³⁾.

Several studies were carried out to see how the pesticide influenced pathogenic fungi. ⁽¹⁴⁾found that the pesticide imidacloprid gave the least inhibitory effect on the fungal growth (11.1%), followed by deltamethrin (36.7%). ⁽¹⁵⁾indicated the compatibility between *Beauveria bassiana* and the pesticide Imidacloprid that enhanced the fungus's efficiency in controlling the whitefly *Bemisia argentifolii*.

The results of exposure of whitefly nymphs to various combinations of control agents (Table 4) showed that the combination of the pesticide Imidacloprid + *Isaria fumosorosea* gave the highest mortality rate in the whitefly nymphs after three days of treatment (79.6%) with a significant difference from the mineral oil And the fungi alone, followed by that mineral oil with *Beauveria bassiana* (78.6%). After seven days of treatment, the superiority of the pesticide Imidacloprid + *Isaria fumosorosea* continued (91.7 %) without significant difference from other combinations, but with significant difference from the agents used alone.

Table (5): The Percentage Mortality of Whitefly Nymphs Due to Exposure to Imidocloprid, Mineral oil, B. BassianaBbk and I. fumosorosea and Their Combinations

Treatment	Dose rate	%Morality	
		After 3 days	After 7 days
Imidocloprid	Half field dose	71.8 ab	73.3 b
Im+ Isf	Half field dose+10 ⁷	79.6 a	91.7 a
Im + Bbk	Half field dose+10 ⁷	75.3 ab	87.6 a
Mineral oil	Half field dose	66.9 b	69.6 bc
Mineral oil+ Isf	Half field dose+10 ⁷	73.0 ab	86.3 a
Mineral oil+ Bbk	Half field dose+10 ⁷	78.6 ab	85.6 a
Isf	10 ⁷	49.6 c	59.3 c
Bbk	10 ⁷	54.6 c	62.3 c

The same letter in column indicated to non significant differences

Combining pesticides with low doses and entomopathogenic fungi can act as a positive action. This is always beneficial as it reduces the doses of the pesticides and thus reduces environmental pollution as well as resistance of insect pests ^{(16),(17)} indicated the compatibility between Beauveriabassiana and the pesticide Imidacloprid and enhanced the fungus's efficiency in controlling the whitefly Bemisiaargentifolii,

The oil mixture enhances the adhesion of the spores to the insect, and later the infection, and the use of oil in compatible concentrations causes a high mortality.⁽¹⁸⁾ On the basis of germination, fungal growth, and spore formation, researchers investigated the compatibility of eight vegetable oils with B. bassiana and M.anisopliae.

Conflict of Interest – Nil

<http://annalsofrscb.ro>

Source of Funding- **Self**

Ethical Clearance – **Not required**

References

1. **Nombela, G. and Muñiz, M.** Host plant resistance for the management of *Bemisiatabaci*: a multi-crop survey with emphasis on tomato. In: Stansly PA, Naranjo SE (eds) *Bemisia*: bionomics and management of a global pest. Springer Science Business Media B.V. 2010., 357–383.
2. **Mascarin, G. M.; Kobori, N. N.; Quintela, E. D. and Delalibera, I.** The virulence of entomopathogenic fungi against *Bemisiatabacibiotype* B (Hemiptera: Aleyrodidae) and their conidial production using solid substrate fermentation. *Biological Control*, 2013. 66(3), 209–218.
3. **Zhang, C.; Ali, S.; Musa, P.D.; Wang, X.M. and Qiu, B.L.** Evaluation of the pathogenicity of *Aschersoniaaleyrodis* on *Bemisiatabaci* in the laboratory and greenhouse. *Biocontrol Sci. Technol.* 2017, 27, 210–221.
4. **Abbott, W. S.** A Method of Computing the Effectiveness of an Insecticide, *Journal of Economic Entomology*. 1925. 18(2): 265–267.
5. **Moorhouse ER, Charnley AK, Gillespie AT.** Review of the biology and control of the vine weevil, *Otiorhynchussulcatus* (Coleoptera: Curculionidae). *Ann. Appl. Biol.* 1992.. 121(2):431-454.
6. **Sain, S.K. ; Monga, D. ; Kumar, R. ; Nagrale, D.T. ; Kranthi, s. and Kranthi, K.R.** Comparative effectiveness of bioassay methods in identifying the most virulent entomopathogenic fungal strains to control *Bemisiatabaci*(Gennadius) (Hemiptera: Aleyrodidae). *Egyptian Journal of Biological Pest Control*, 2019 .29:31.
7. **Quesada-Moraga, E.; Maranhao ,E.A.A.; Valverde- García, P. and Santiago-Álvarez, C.** Selection of *Beauveria bassiana* isolates for control of the whiteflies *Bemisiatabaci* and *Trialeurodes vaporariorum* on the basis of their virulence, thermal requirements, and toxicogenic activity. *Biological Control*. 2006. 36:274-287.
8. **Rahim, E.; Ahmad, S.S.; Dzolkhifli, O. and Nur, A.A.** First record of *Isaria fumosorosea* Wize (Deuteromycotina: Hyphomycetes) infecting *Bemisiatabaci*(Gennadius) (Hemiptera: Aleyrodidae) in Malaysia. *J. Entomol.* 2013, 10, 182–190. **Kereselidze, M.; Draganova, S.; Pilarska, D. and Linde, A.** Susceptibility of *Lymantria monacha* and *L. dispar* to the entomopathogenic fungus *Isaria fumosorosea* Wize. *Sumar*. 2014, 138, 185–191.
9. **Vinoth, B.K.; Kuttalam, S. and Chandrasekaran, S.** Efficacy of a new insecticide spirotetramat against cotton whitefly. 2009. *Pestic. Res. J.*, 21: 45 – 48.
10. **Vinoth, B.K.; Kuttalam, S. and Chandrasekaran, S.** Efficacy of a new insecticide spirotetramat against cotton whitefly. 2009. *Pestic. Res. J.*, 21: 45 – 48.

- 11.Amutha, M.; Banu, J. G.; Surulivelu, T. and Gopalakrishnan, N.** Effect of commonly used insecticides on the growth of white Muscardine fungus, *Beauveria bassiana* under laboratory conditions. Journal of Biopesticides, 2010. 3(1): 143-1.
- 12.Amutha, M.; Banu, J. G.; Surulivelu, T. and Gopalakrishnan, N.** Effect of commonly used insecticides on the growth of white Muscardine fungus, *Beauveria bassiana* under laboratory conditions. Journal of Biopesticides, 2010. 3(1): 143-1.
- 13.Cuthbertson, A.G.S.; Mathers, J.J.; Northing, P.; Luo WeiQi and Walters, K.F.A.** The susceptibility of immature stages of *Bemisia tabaci* to infection by the entomopathogenic nematode *Steinernema carpocapsae*. Russian Journal of Nematology, 2007. 15(2):153-156.
- 14.Mascarin, G. M. ;Kobori, N.N. ; Quintela, E.D. ; Arthurs, S.P. and Ju´nior. I. D.** Toxicity of non-ionic surfactants and interactions with fungal entomopathogens toward *Bemisia tabaci* biotype B. BioControl, 2014. 59:111–123.
- 15.Zou, C.; Li, L.; Dong, T.; Zhang, B. and Hu, Q.** Joint action of the entomopathogenic fungus *Isaria fumosorosea* and four chemical insecticides against the whitefly *Bemisia tabaci*. 2014. Biocontrol Sci. Technol., 24, 315–324.
- 16.Bateman, T. S., and Crant, J. M.** The proactive component of organizational behavior. Journal of Organizational Behavior, 1993. 14, 103-118.
- 18.Ummidi, V. R. S., and Vadlamani, P.** Preparation and use of oil formulations of *Beauveria bassiana* and *Metarhiziumanisopliae* against *Spodopteralitura* larvae. African Journal of Microbiology Research, 2014. 8(15), 1638-1644.