

A Descriptive Study to Assess Obesity and Selected Food Habits among Adolescents at Pune

Jasneet Kaur

Associate Professor, Symbiosis College of Nursing, Symbiosis International (Deemed University)

ABSTRACT

Obesity is a problem in which overload fat usually accumulate in the body and leads to unfavourable effect on health of the person which further leads to decreased life expectancy. The purpose of the study is to assess obesity and selected food habits among adolescents with a view to provide data regarding the prevalence of obesity. The descriptive approach was considered appropriate for the present study as it is aimed to assess the obesity and selected food habits among adolescents. Non-Experimental Research Design is used in the present study. 50 participants were selected with the help of purposive sampling technique. The tool was prepared to identify obesity and selected food habits. The results showed that maximum adolescents preferred junk food. More than 28% of adolescents preferred canteen food. 28% of adolescents was unhealthy out of that 22% were males. It is concluded and recommended that obesity is dreadful condition and hence to be take care at early stage only. Therefore awareness programs should be executed to prevent this.

Keywords

Obesity, Food Habits, Adolescents

INTRODUCTION

Obesity is a problem in which overload fat usually accumulate in the body and leads to unfavourable effect on health of the person which further leads to decreased life expectancy. Body Mass Index is one of the method which compares weight and height. It concludes the level of obesity as overweight if ranges between 25 and 30 kg/ m², and obese if more than 30. The overweight People are increasing around the world compared to the underweight. Developing countries have also coupled in the line along with the developed countries in the race of obesity. It is growing in all developing areas even in the countries plagued by hunger. As per WHO, Prevalence of childhood obesity across the globe showed more than 2 percent in developed and 10% greater in developing in preschool children. India has the rate of 1% in a preschool child obesity.

Obesity can occur at any age. It usually happens to be increased with age. Infants with excessive weight gain have an increased incidence of obesity in later life. Childhood obesity usually resulted as interaction between hereditary and environmental factors. This is the need of today to explore more the prevalence, risk factors and deleterious effects of obesity.

Problem Statement

A descriptive study to assess obesity and selected food habits among adolescents at Pune

Purpose of the Study

The purpose of the study is to assess obesity and selected food habits among adolescents with a view to provide data regarding the prevalence of obesity.

Objectives of the Study

- To identify the prevalence of obesity among adolescents
- To assess food habits among adolescents.
- To determine the association of obesity among adolescents with socio-demographic variables.

METHODOLOGY

The **descriptive approach** was considered appropriate for the present study as it is aimed to assess the obesity and selected food habits among adolescents. Non-Experimental Research Design is used in the present study. 50 participants were selected with the help of purposive sampling technique. The tool was prepared to identify obesity and selected food habits. Analysis of data was done in accordance with the objectives of the study. Study was done by using the descriptive and inferential statistics as calculated by frequency, percentage, mean score, standard deviation and Karl Pearson's correlation, chi-square test, BMI of school students assessed as per percentiles in order to assess the obesity and selected food habits.

RESULTS

Section-I: Sample Characteristics

Majority (84%) were from the age of 11 -12 years followed by the age of 12-13yrs (2%). Majority (78%) of students were males and females were (32%). Most of Students were from Urban areas (88%) and only (12%) were from Rural areas. Regarding their religion about (82%) were from Hindu religion. Half (52%) of students were from nuclear family and (48%) were from joint families.

Majority (70%) of students have 2 siblings followed by (22%) subjects followed by (22%) subjects having more than 2 siblings and only (8%) have one sibling.

Section-II

Table 1: Prevalence of Obesity among School Students

S. No.	BMI	Percentile	n	%	Age
1.	Under Weight	> 5 th	5	10	
2.	Healthy Weight	5 th upto 85 th	34	68	

3.	Over Weight	85 th to < 95 th	7	14
4.	Obese	= or > 95 th	4	8

The above table represents that 68% of the students whose percentile is 5th upto 85th are having the BMI of healthy weight and only 8% of the students whose percentile is equal to or more than 95th, are being BMI of obese category.

Section III - Food Habits among School Students

Table 2: Fad Diet Scale

N=150				
Sr. No.	Canteen Items	Never n (%)	Sometimes n (%)	Always n (%)
1.	Biscuits	9 (18%)	33(66%)	8(16%)
2.	Chocolate	6 (12%)	32(64%)	12(24%)
3.	Toffees	4(8%)	34(68%)	12(24%)
4.	Weffers	28(56%)	18(36%)	14(8%)
5.	Kurkure	7(14%)	30(60%)	13(26%)
6.	Chips	5(10%)	33(66%)	12(24%)
7.	Fun pops	11(22%)	29(58%)	10(20%)
8.	Bhel-puri	20(40%)	26(52%)	4(8%)
9.	Samosa	10(20%)	34(68%)	6(12%)
10.	Spring roll	17(34%)	29(58%)	4(8%)
11.	Patty	11(22%)	36(72%)	3(6%)
12.	Ice gola	17(34%)	25(50%)	8(16%)
13.	Icecream	9(18%)	27(54%)	14(28%)
14.	Tea	18(36%)	10(20%)	22(44%)
15.	Cold drinks	7(14%)	31(62%)	12(24%)

The table represents that maximum no. of students (66%) sometimes eat biscuit and only (18%) are the students who have never ingested the biscuits and mere (16%) of the students are always in the habit of eating biscuits.

The table depicts that maximum of students (60%) and (66%) sometimes eats kurkure and chips respectively. Only (14%) and (10%) are the students who have never ingested the same respectively. Rest (26%) and (24%) of the students are always in the habits of eating kurkure and chips respectively.

The table assays that maximum (52%) of students sometimes eat bhel-puri and (40%) are the students who have never ingested bhel-puri and mere (8%) of the students are always in the habit of eating bhel-puri.

The table shows that maximum number of students 34(68%) respectively sometimes eat samosa's. Only (20%) of students have never ingested same. Rest (12%) always eat the same.

The table depicts that maximum (58%) of students sometimes eat spring roll while (34%) never eat the same rest (8%) of students always eat the same.

The table represents that maximum (72%) of students sometimes eat patty and only (22%) and (6%) of students never and always eat the same respectively.

The table shows that maximum (44%) of students are always in the habit of having tea. Only (36%) are the students who never taken tea. Rest (20%) of the students sometimes take the same.

Table 3: Times of Food taken in a Day by Adolescents

S. No.	Time of Food Taken	Frequency (n)	Percentage (%)
1.	Three times only	32	64 %
2.	More than three times	6	12 %
3.	Less than three times	13	24 %

Table 3 shows more than half (64%) students take food three times a day followed by (24%) students who take food less than three times in a day and (12%) students who take more than three times food in a day.

Table 4: Type of Lunch Preferred

S. No.	Type of Lunch Preferred	Frequency (n)	Percentage (%)
1.	Homemade	36	72 %
2.	Canteen snacks	14	28 %

Table 4 shows that about 3/4th (72%) of students prefer homemade lunch in school and 1/4th (28%) students prefer canteen snacks in school.

Table 5: Means of Transport used by School Students for Coming to School

S. No	Means of Transport Used	Frequency (n)	Percentage (%)
1.	Walking	2	4%
2.	Bicycle	6	12 %
3.	Private Vehicle	42	84 %

Table 5 shows that more than 3/4th (84%) students come to school private vehicle (car, bike) and rest (12%) and (4%) come by bicycle and by walking.

Section IV: Association of Obesity and Socio Demographic Variables

Maximum (84% are from age group 11-12 yrs from which 56% students have normal weight but still 28% students are unhealthy. More than 3/4th (78%) students are males out of which 48% students have normal weight and rest of 22% are unhealthy. About 88% students are from urban areas out of which 58% students have normal weight and rest are unhealthy. About 48% of students are from joint family from which 30% have normal weight and rest of 18% are unhealthy. More than 2/4th (52%) students are from Nuclear family from which 38% students have normal weight and rest of 14% are unhealthy. Maximum (70%) of students have two siblings out of which 48% students have normal weight and rest of 22% students are unhealthy.

CONCLUSION

Children are the one who are very much attracted towards the foods available in school. They are not aware that these foods can cause obesity. They are only concerned about their taste. Bad food habits lead to assess the effect of food habits on obesity.

REFERENCES

- [1] Kotani K, Nishida M, Yamashita S, Funaharhi T, Fujoka S, Ishikawa K, Tarwi S and Matsuzawa Y, Two decades of annual medical examination in Japanese obese children, *Internet J of Obes and Rel Metabolic Disorder*, 21: 912-921, 1997.
- [2] Gurney M, Gorstein J, The global prevalence of obesity - an initial overview of the available data, *World Health Stats Quarterly*, 41: 251-254, 1988.
- [3] Kaur S, Kapil U, Singh P (2005) Patterns of chronic diseases amongst adolescent obese children in developing countries. *Curr. Sci* 88: 1052-1056.
- [4] Sawane, K., & Barde, S. (2019). Mother's knowledge on nutrition and incidence of malnutrition. *Indian Journal of Public Health Research and Development*, 10(1), 32-34. doi:10.5958/0976-5506.2019.00008.1
- [5] James J, Kerr D (2005) "Prevention of childhood obesity by reducing soft drinks", *Int. J. Obes (Lond)* 29 (Suppl 2) : 854-7.
- [6] Horton TJ, Drougas H, Brochey A, Reed GW, Peters JC, Hill JO (1995) "Fat and CHO overfeeding in humans: different effects on energy storage", *Am. J. Clin. Nutr.* 62(1): 19-29.
- [7] North American Society for Pediatric Gastroenterology, Hepatolog and Nutrition (PDF).
- [8] Brijalska IJ, Kumar S, Stewart PM (1997) Does central obesity reflect "Cushing's disease of the omentum?" *Lancet* 349 (9060): 1210-3
- [9] Must A, Jacques PF, Dallal GE Bajema CJ, Dietz WH (November 1992). "Long term morbidity and mortality of overweight adolescents. A follow-up of the Harward growth study of 1922-1935", *The new England journal of medicine* 327 (19): 1350-5.

- [10] Strychar J (January 2006) "Diet in the management of weight loss" *CMAJ* 174(1): 56-63.
- [11] Epstein LH, Roemmich JN, Robinson Jt, (2008).
- [12] Ong KK, Ahmed ML, Emmett PN, Preece MA, Bunger DB (2000) "Association between postnatal catch up growth and obesity in childhood: prospective cohort study", *BMJ* 320.
- [13] James PT, Leach R, Kalamara E &Shayeghi M (2001), The worldwide obesity epidemic. *Obesity Research*, 9 (Suppl 4) 228 S - 233 S.
- [14] Green V, (2005), The domino effect: obesity, type 2 diabetes and cardiovascular disease, *British Journal of Community Nursing*, 10, 358-361.
- [15] Wabitsch M, (2000), Overweight and obesity in European children: Definition and diagnostic procedure, risk factors and consequences for later health outcome. *European Journal of pediatrics*, 159 (Suppl.1), S8-S13, doi: 10.1007/PL00014368.
- [16] Vaska V and Volkmer R, (2004), increasing prevalence of obesity in South Australian 4-year-olds: 1995-2002, *Journal of pediatrics and child health*, 40, 353-355, doi: 10.1111/j-1034-4810.2004.00401.x.
- [17] Janssen I, Katzmarzyk PT Boyce WF King MA and Pickett W(2004), Overweight and obesity in Canadian adolescents and their associations with dietary in habits and physical activity patterns, *Journal of Adolescent Health*, 35 (360-367).