

## **Evaluation of Preference of Root Canal Irrigants by Students for Pulpectomy**

**Aniruddh Menon**

Saveetha Dental College and Hospital,  
Saveetha Institute of Medical and Technical Sciences,  
Saveetha University, Chennai-77  
151501006.sdc@saveetha.com

**Mebin George Mathew**

Senior Lecturer ,  
Department of Pediatric and Preventive Dentistry,  
Saveetha Dental College and Hospital,  
Saveetha Institute of Medical and Technical Sciences,  
Saveetha University, Chennai-77  
mebingeorgem.sdc@saveetha.com

**Jayanth Kumar Vadivel**

Reader  
Department of Oral Medicine and Radiology,  
Saveetha Dental College and Hospital,  
Saveetha Institute of Medical and Technical Sciences,  
Saveetha University, Chennai-77  
jayanthkumar@saveetha.com

**Corresponding Author:**

**Mebin George Mathew**

Senior Lecturer ,  
Department of Pediatric and Preventive Dentistry,  
Saveetha Dental College and Hospital,  
Saveetha Institute of Medical and Technical Sciences,  
Saveetha University, Chennai-77  
mebingeorgem.sdc@saveetha.com

### **ABSTRACT:**

The present study was undertaken to evaluate the preference of root canal irrigants for pulpectomy of primary teeth in at Saveetha Dental college. Retrospective data of 2438 patients was obtained and segregated who visited between June 2019 to April 2020. Once the data was obtained it was statistically analyzed using SPSS by IBM version 20. The most commonly used out of the irrigants preferred in the present study was saline (59%) followed by saline with a combination of EDTA (32.5%), followed by EDTA alone (8.4%) and the least used irrigant was a combination of saline and sodium hypochlorite (0.1%). Saline was the most preferred irrigant in both the undergraduate and postgraduate clinics. Within the limits of the present study it was observed that the most common type of irrigant that was used is saline (59.03%). Further studies are to be done to formulate a clear and distinct clinical practice guideline for

pulpectomy procedure. Further awareness programmes that should be conducted to bring out better clinical outcomes and help the society holistically.

**Key words:** EDTA; irrigant; primary; pulpectomy; saline.

## INTRODUCTION:

There are various situations (Christabel and Linda Christabel, 2015; Gurunathan and Shanmugaavel, 2016; Packiri, Gurunathan and Selvarasu, 2017) arising in children that require intervention by practitioners (Govindaraju, Jeevanandan and Subramanian, 2017b; Packiri, Gurunathan and Selvarasu, 2017; Ravikumar, Jeevanandan and Subramanian, 2017). With the advent of time, there have been various trends in the occurrence of caries in pediatric patients (Subramanyam *et al.*, 2018) and various approaches to prevent and treat it (Somasundaram, 2015; Govindaraju, 2017b; 'Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children - Review', 2018).

A pulpectomy procedure may be performed on primary teeth when the coronal portion of the pulp tissue and the tissue entering the pulp canals are vital but show clinical evidence of hyperemia or if the root canals show evidence of necrosis or suppuration. It is unwise to maintain or retain untreated infected primary teeth in the oral cavity. They may be opened for drainage of the affected tissue and often remain asymptomatic for an indefinite period. However, they are a source of infection and should be treated or removed. The morphology of the root canals in primary teeth makes endodontic treatment difficult. Mature first primary molar canals are often so small that they are inaccessible even to the smallest barbed broach. If the canal cannot be properly cleansed of necrotic material, sterilized, and adequately filled, endodontic therapy is more likely to fail. The success of endodontic therapy in primary teeth strongly depends on achieving an adequate level of disinfection within the root canals. Mechanical instrumentation alone is unlikely to be sufficient in attaining such disinfection, considering the root resorption process and the complex anatomy of the root canal system, characteristic of primary molars, and the risk of damage to the permanent germ (Kargül, Tanboga and Altinok, 2010). There are various exclusive file systems have been recently developed for the purpose of performing pulpectomy in primary teeth and their efficacy has been tested with time (Jeevanandan, 2017; Jeevanandan and Govindaraju, 2018a; Panchal, Jeevanandan and Subramanian, 2019). Evidence has shown that the mechanical instrumentation technique with files is limited because it tends to leave significant portions of the infected canal walls untouched; thus, a great number of viable pathogenic microorganisms persist, lodged together with dentin debris and necrotic pulp-tissue remnants inside the dentin tubules, canal ramifications, and resorption craters (Gondim *et al.*, 2012). The major reduction of bacteria in the root canals is achieved by the mechanical action of endodontic files and by irrigation. The need for medication increases in those cases where infection resists regular treatments and the therapy cannot be successfully completed owing to the presence of pain or continuing exudation. Until the mid-1980s, there was a preference for using strong phenolic intracanal antiseptics such as formocresol, camphorated paramonochlorophenol (CPMC), cresatin etc. CMCP proved to be one of the most toxic and irritating phenolic antiseptic followed by cresatin, formocresol and camphorated phenol (CP) (Athanassiadis, Abbott and Walsh, 2007) (Masillamoni, Kettering and Torabinejad, 1981). Therefore, it is necessary to significantly reduce or to eradicate, to the extent possible, the

microorganisms and their by-products present to the pulp canals by employing clinically effective and biocompatible irrigants, which also aid in dissolving organic debris (Ito, Junior and Paula-silva, 2011).

Irrigation currently represents the best method in paediatric pulpectomy for the lubrication and flushing away of loose necrotic and contaminated materials during instrumentation (Ito, Junior and Paula-silva, 2011; Kauret *et al.*, 2014). In clinical practice, different intracanal irrigants have been proposed for primary teeth, such as sodium hypochlorite (NaOCl), chlorhexidine gluconate, ethylenediaminetetraacetic acid (EDTA), citric acid and hydrogen peroxide (Kauret *et al.*, 2014), causing confusion among practitioners especially those who are less experienced. Thus, it is a challenge for dentists to choose the most appropriate irrigant agent when performing pulp canal treatments. Although several studies have tried to establish the effectiveness of diverse disinfecting substances, the irrigating solution that could be considered as the ideal agent irrigating agent has still not been found (Önçağ, Hoşgör and Hilmioğlu, 2003; Vasconcelos, Luna-Cruz and De-Deus, 2007; Tulsani, Chikkanarasaiah and Bethur, 2014). All of this is of utmost importance as it will influence the end product of obturation (Govindaraju, 2017a; Govindaraju, Jeevanandan and Subramanian, 2017a) as well as patient care (Nair *et al.*, 2018). Our team has rich experience in research and we have collaborated with numerous authors over various topics in the past decade (Ariga *et al.*, 2018; Basha, Ganapathy and Venugopalan, 2018; Hannah *et al.*, 2018; Hussain *et al.*, 2018; Jeevanandan and Govindaraju, 2018b; Kannan and Venugopalan, 2018; Kumar and Antony, 2018; Manohar and Sharma, 2018; Menon *et al.*, 2018; Nandakumar and Nasim, 2018; Nandhini, Babu and Mohanraj, 2018; Ravinthar and Jayalakshmi, 2018; Seppan *et al.*, 2018; Teja, Ramesh and Priya, 2018; Duraisamy *et al.*, 2019; Gheena and Ezhilarasan, 2019; Hema Shree *et al.*, 2019; Rajakeerthi and Ms, 2019; Rajendran *et al.*, 2019; Sekar *et al.*, 2019; Sharma *et al.*, 2019; Siddique *et al.*, 2019; Janani, Palanivelu and Sandhya, 2020; Johnson *et al.*, 2020; Jose, Ajitha and Subbaiyan, 2020).

The aim of the present study was to evaluate the preference of root canal irrigants for pulpectomy of primary teeth in a university setting.

## **MATERIALS AND METHODS:**

The present study involved a total of 2438 patients that underwent single visit pulpectomy procedures. The irrigants studied included saline, EDTA, Saline + EDTA, saline + sodium hypochlorite. The study was performed in a university setting at Saveetha Dental College and Hospitals. Thus the data obtained from the patients is of the same geographic location and ethnicity. The ethical approval for collection of retrospective data from the dental patient management archives was obtained from the Institutional Ethics Board. (IRB Approval No: SIHEC/2020/DIASDATA/0619-0320) The period of the study was between 1<sup>st</sup> June 2019 to 1<sup>st</sup> April 2020. Once the data was collected the same was verified by using photographs by two external reviewers who were blinded on the hypothesis from the present study. This was done to eliminate the chances of sampling bias. Before the commencement of the study a clear well defined inclusion criteria was defined. The inclusion criteria included that:

- Patient has been treated by a resident of Saveetha Dental College, either an undergraduate or postgraduate student.

- Should have undergone single visit pulpectomy
- Should have been within the age group of 1-12 years. Patients were segregated into three age groups; Group 1- 1 - 4 years ; Group 2- 5 to 8 years; Group 3- 9-12 years.

Out of the study population that was chosen for the study there was no segregation process, as this would result in sampling bias. The data segregation was done according to various parameters such as speciality of clinic in which patient was treated, age of the patient, gender of the patient etc.

The data that was then tabulated was reviewed by an external reviewer and screened for internal validity of the study. The data was then exported to SPSS Software by IBM Version 20 for Statistical Analysis. Descriptive statistics was performed followed by Correlation tests to see any kind of correlation or Association between the different variables taken in the present study. Descriptive statistics and Chi square test was used to determine the correlation between the variables where P value < 0.05 is considered statistically significant with a confidence interval of 95%.

## RESULTS:

Of the 2438 cases that underwent pulpectomy treatment and were included as part of the study, 1421 cases ( 58.3%) were males and 1017 cases were females (41.7%). The patients in the present study were within the age group of 1-12 years with a mean age of  $4.8 \pm 1.7$  years. Majority of the patients were between the ages of 1-6 years (81.9%) and the remaining between 6-12 years (19.1%)

The teeth most frequently to have undergone pulpectomy were 84,85 (24.9%) followed by 74,75 (23.4%) . 2215 of the procedures were performed in the postgraduate clinic (90.9%), 200 (8.2%) in undergraduate clinics and 23 (0.9%) by specialists.

Out of the four irrigants in the present study the most commonly used was saline (59%) followed by saline with a combination of EDTA (32.5%), followed by EDTA alone (8.4%) and the least used irrigant was a combination of saline and sodium hypochlorite (0.1%). Saline was the most preferred irrigant in both the undergraduate and postgraduate clinics, and sodium hypochlorite was used only in the postgraduate clinics. A high correlation and association was noted between age of the patient and the type of irrigant used ( $p=0.002$ )

## DISCUSSION:

In recent times, commonly used irrigation solutions include NaOCl, EDTA, citric acid, chlorhexidine (CHX), and electrochemical activation (ECA) of different solutions. However, the apparent toxicity of NaOCl and the effects of the chlorates ( $\text{ClO}_3^-$  and  $\text{ClO}_2^-$ ) should be recognized . In humans, these chemical substances are the products of oxidative damage to red blood cells, hemolytic anemia and methemoglobin formation and are also cytotoxic for fibroblasts.(Paudelet *et al.*, 2011) These should be kept in mind before choosing an irrigant.

Currently, there is no agreement among paediatric dentists concerning the best irrigant solution for use against pulp pathogens involved in irreversibly inflamed/infected or non-vital primary teeth; this absence of consensus is probably due to the lack of evidence based research research to prove the superiority of any irrigants(Gondimet *et al.*, 2012) agent . In their narrative review of

intracanal(Kaure *et al.*, 2014)irrigants for primary teeth, Kaur *et al.* proposed six requirements that the ideal irrigant must possess: (1) broad antimicrobial spectrum, (2) high efficacy against anaerobic and facultative microorganisms organized in bio- films, (3) ability to dissolve necrotic pulp-tissue remnants, (4) ability to inactivate endotoxin, (5) ability to prevent the formation of or to dissolve the smear layer during instrumentation, and (6) a non-toxic (to periodontal tissues), non-caustic nature that does not cause an allergic reaction. Although there is a lack of consensus among various irrigants that have been included in the study, the most frequently used irrigant was saline. However saline does not satisfy most of the features that are included as part of the ideal criteria that is required for an ideal root canal irrigant. The advantage of saline is that it is non toxic to the periapical and periradicular tissues as reported by Turkun *et al* (Türkün, Gökay and Özdemir, 1998) and Yesilsoy *et al* (Yesilsoy *et al.*, 1995). It is evident from the study carried out by Yoshida *et al* (Yesilsoy *et al.*, 1995; Yoshida *et al.*, 1995) where antimicrobial activity of various irrigants were tested, saline has the least antimicrobial activity. Another factor that has to be considered is that the present study was carried out in a university setting where all participants were in a state of learning which could lead to accidents using irrigants such as NaOCl(Mehdipouret *al.*, 2007)Thus, to prevent the same from occurring most students would have preferred saline and with time might progress to other irrigants. The preference for EDTA and combination of EDTA and saline as irrigants was seen to be higher with post graduate students. This could be due to the fact that most postgraduates would prefer rotary instruments compared to undergraduates.(Jeevanandan and Govindaraju, 2018a)Our institution is passionate about high quality evidence based research and has excelled in various fields ( (Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; VijayashreePriyadharsini, SmilineGirija and Paramasivam, 2018; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Ramadurai *et al.*, 2019; Sridharan *et al.*, 2019; VijayashreePriyadharsini, 2019; Chandrasekaret *al.*, 2020; Mathew *et al.*, 2020; R *et al.*, 2020; Samuel, 2021)

The limitations of the present study include that all irrigants available have not been included in the study, all the participants are students and practitioners can be involved to obtain a more concise knowledge and approach towards the same.

## CONCLUSION:

Within the limits of the present study it is observed that the most common type of irrigant that is used is saline (59.03%) and the least commonly used is the combination of saline and hypochlorite(0.45%). Further studies are to be done to formulate a clear and distinct clinical practice guideline for pulpectomyprocedure.Further awareness programmes that should be conducted to bring out better clinical outcomes and help the society holistically.

## AUTHOR CONTRIBUTIONS:

AniruddhMenon carried out the retrospective study, planning the study design, collection and analysis of data and drafted the manuscript. Nashra Kareem and Jayanth Kumar Vadivel aided in conception of the topic, supervision and appraisal of the manuscript.

## ACKNOWLEDGEMENTS:

We thank Saveetha Dental College and Hospitals, Chennai for access to the retrospective data

## CONFLICT OF INTEREST:

No conflict of Interest.

## REFERENCES:

1. Ariga, P. *et al.* (2018) 'Determination of correlation of width of Maxillary Anterior Teeth using Extraoral and Intraoral Factors in Indian Population: A systematic review', *World journal of dentistry*, 9(1), pp. 68–75.
2. Athanassiadis, B., Abbott, P. V. and Walsh, L. J. (2007) 'The use of calcium hydroxide, antibiotics and biocides as antimicrobial medicaments in endodontics', *Australian dental journal*, 52(1 Suppl), pp. S64–82.
3. Basha, F. Y. S., Ganapathy, D. and Venugopalan, S. (2018) 'Oral hygiene status among pregnant women', *Journal of advanced pharmaceutical technology & research*, 11(7), p. 3099.
4. Chandrasekar, R. *et al.* (2020) 'Development and validation of a formula for objective assessment of cervical vertebral bone age', *Progress in orthodontics*, 21(1), p. 38.
5. Christabel, S. L. and Linda Christabel, S. (2015) 'Prevalence of Type of Frenal Attachment and Morphology of Frenum in Children, Chennai, Tamil Nadu', *World Journal of Dentistry*, pp. 203–207. doi: 10.5005/jp-journals-10015-1343.
6. Duraisamy, R. *et al.* (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295.
7. Ezhilarasan, D., Apoorva, V. S. and Ashok Vardhan, N. (2019) 'Syzygiumcumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121.
8. 'Fluoride, Fluoridated Toothpaste Efficacy And Its Safety In Children - Review' (2018) *International Journal of Pharmaceutical Research*. doi: 10.31838/ijpr/2018.10.04.017.
9. Gheena, S. and Ezhilarasan, D. (2019) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & experimental toxicology*, 38(6), pp. 694–702.
10. Gondim, J. O. *et al.* (2012) 'Effect of a calcium hydroxide/chlorhexidine paste as intracanal dressing in human primary teeth with necrotic pulp against Porphyromonasgingivalis and Enterococcus faecalis', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 22(2), pp. 116–124.
11. Govindaraju, L. (2017a) 'Clinical Evaluation of Quality of Obturation and Instrumentation Time using Two Modified Rotary File Systems with Manual Instrumentation in Primary Teeth', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2017/30069.10602.
12. Govindaraju, L. (2017b) 'Effectiveness of Chewable Tooth Brush in Children-A Prospective Clinical Study', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2017/24238.9528.
13. Govindaraju, L., Jeevanandan, G. and Subramanian, E. M. G. (2017a) 'Comparison of quality of obturation and instrumentation time using hand files and two rotary file systems in primary molars: A single-blinded randomized controlled trial', *European Journal of Dentistry*, pp. 376–379. doi: 10.4103/ejd.ejd\_345\_16.
14. Govindaraju, L., Jeevanandan, G. and Subramanian, E. M. G. (2017b) 'Knowledge and practice of rotary instrumentation in primary teeth among indian dentists: A questionnaire


- survey', *Journal of International Oral Health*, p. 45. doi: 10.4103/jioh.jioh\_4\_17.
15. Gurunathan, D. and Shanmugaavel, A. (2016) 'Dental neglect among children in Chennai', *Journal of Indian Society of Pedodontics and Preventive Dentistry*, p. 364. doi: 10.4103/0970-4388.191420.
16. Hannah, R. *et al.* (2018) 'Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens', *Journal of advanced pharmaceutical technology & research*, 11(3), p. 1012.
17. Hema Shree, K. *et al.* (2019) 'Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma - a Systematic Review with Meta Analysis', *Pathology oncology research: POR*, 25(2), pp. 447–453.
18. Hussainy, S. N. *et al.* (2018) 'Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up', *Journal of conservative dentistry: JCD*, 21(5), pp. 510–515.
19. Ito, I. Y., Junior, F. M. and Paula-silva, F. (2011) 'Microbial culture and checkerboard DNA–DNA hybridization assessment of bacteria in root canals of primary teeth pre-and post-endodontic therapy with a calcium ...', *journal of paediatric ....* Available at: [https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-263X.2011.01131.x?casa\\_token=MKWyqU2I6p8AAAAA:6Lp5XmXoXa9iUUaKOjJ\\_kn6qozaNRbKodRLAA5Igg87C9GyGHuraP0tQ86CPuHguyASbz\\_\\_kovPgkec](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-263X.2011.01131.x?casa_token=MKWyqU2I6p8AAAAA:6Lp5XmXoXa9iUUaKOjJ_kn6qozaNRbKodRLAA5Igg87C9GyGHuraP0tQ86CPuHguyASbz__kovPgkec).
20. Janani, K., Palanivelu, A. and Sandhya, R. (2020) 'Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality: an in vivo study', *Brazilian dental science*, 23(1). doi: 10.14295/bds.2020.v23i1.1805.
21. Jeevanandan, G. (2017) 'Kedo-S Paediatric Rotary Files for Root Canal Preparation in Primary Teeth – Case Report', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2017/25856.9508.
22. Jeevanandan, G. and Govindaraju, L. (2018a) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European Archives of Paediatric Dentistry*, pp. 273–278. doi: 10.1007/s40368-018-0356-6.
23. Jeevanandan, G. and Govindaraju, L. (2018b) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 19(4), pp. 273–278.
24. Johnson, J. *et al.* (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH', *Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362.
25. Jose, J., Ajitha and Subbaiyan, H. (2020) 'Different treatment modalities followed by dental practitioners for Ellis class 2 fracture – A questionnaire-based survey', *The open dentistry journal*, 14(1), pp. 59–65.
26. Kannan, A. and Venugopalan, S. (2018) 'A systematic review on the effect of use of impregnated retraction cords on gingiva', *Journal of advanced pharmaceutical technology & research*, 11(5), p. 2121.
27. Kargül, B., Tanboga, I. and Altinok, B. (2010) 'Conventional endodontic treatment of

- primary molars using metronidazole as an intra-canal medicament: A pilot study', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 11(4), pp. 196–200.
28. Kaur, R. *et al.* (2014) 'Irrigating solutions in pediatric dentistry: Literature review and update', *J Adv Med Dent Sci Res*, 2, pp. 104–115.
29. Kumar, D. and Antony, S. D. P. (2018) 'Calcified canal and negotiation-A review', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3727.
30. Manohar, M. P. and Sharma, S. (2018) 'A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(6), pp. 716–720.
31. Masillamoni, C. R., Kettering, J. D. and Torabinejad, M. (1981) 'The biocompatibility of some root canal medicaments and irrigants', *International endodontic journal*, 14(2), pp. 115–120.
32. Mathew, M. G. *et al.* (2020) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: Randomized controlled trial', *Clinical oral investigations*, pp. 1–6.
33. Mehdi-pour, O. *et al.* (2007) 'Anatomy of sodium hypochlorite accidents', *choice*, 5(8), p. 9.
34. Menon, S. *et al.* (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', *Colloids and surfaces. B, Biointerfaces*, 170, pp. 280–292.
35. Nair, M. *et al.* (2018) 'Comparative evaluation of post-operative pain after pulpectomy with k-files, kedo-s files and mtwo files in deciduous molars -a randomized clinical trial', *Brazilian Dental Science*, p. 411. doi: 10.14295/bds.2018.v21i4.1617.
36. Nandakumar, M. and Nasim, I. (2018) 'Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis', *Journal of conservative dentistry: JCD*, 21(5), pp. 516–520.
37. Nandhini, J. S. T., Babu, K. Y. and Mohanraj, K. G. (2018) 'Size, shape, prominence and localization of gerdy's tubercle in dry human tibial bones', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3604.
38. Öncü, Ö., Hoşgör, M. and Hilmioğlu, S. (2003) 'Comparison of antibacterial and toxic effects of various root canal irrigants', *International*. Available at: [https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2591.2003.00673.x?casa\\_token=WiwnNxd6I40AAAAA:CAgZKvv2TtxkFlfBGGu\\_BVjceHddeWICGKsaEIIUII9ZDM8Dd\\_7np9BdycuGDDEToXAPIfd0z7-DKF38](https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2591.2003.00673.x?casa_token=WiwnNxd6I40AAAAA:CAgZKvv2TtxkFlfBGGu_BVjceHddeWICGKsaEIIUII9ZDM8Dd_7np9BdycuGDDEToXAPIfd0z7-DKF38).
39. Packiri, S., Gurunathan, D. and Selvarasu, K. (2017) 'Management of paediatric oral ranula: a systematic review', *Journal of clinical and diagnostic research: JCDR*, 11(9), p. ZE06.
40. Panchal, V., Jeevanandan, G. and Subramanian, E. M. G. (2019) 'Comparison of instrumentation time and obturation quality between hand K-file, H-files, and rotary Kedo-S in root canal treatment of primary teeth: A randomized controlled trial', *Journal of Indian Society of Pedodontics and Preventive Dentistry*, p. 75. doi: 10.4103/jisppd.jisppd\_72\_18.
41. Paudel, K. R. *et al.* (2011) 'Different pharmacological solutions in intracanal irrigation', *Nepal Medical College journal: NMCJ*, 13(2), pp. 111–114.




42. Pc, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research*. Available at: <https://europepmc.org/article/med/29624863>.
43. Rajakeerthi and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 22(2), pp. 249–256.
44. Rajendran, R. *et al.* (2019) 'Comparative evaluation of remineralizing potential of a paste containing bioactive glass and a topical cream containing casein phosphopeptide-amorphous calcium phosphate: An in vitro study', *Pesquisabrasileira em Odontopediatria e Clínica Integrada*, 19(1), pp. 1–10.
45. Ramadurai, N. *et al.* (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial', *Clinical oral investigations*, 23(9), pp. 3543–3550.
46. Ramesh, A. *et al.* (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248.
47. Ravikumar, D., Jeevanandan, G. and Subramanian, E. M. G. (2017) 'Evaluation of knowledge among general dentists in treatment of traumatic injuries in primary teeth: A cross-sectional questionnaire study', *European Journal of Dentistry*, pp. 232–237. doi: 10.4103/ejd.ejd\_357\_16.
48. Ravinthar, K. and Jayalakshmi (2018) 'Recent advancements in laminates and veneers in dentistry', *Journal of advanced pharmaceutical technology & research*, 11(2), p. 785.
49. R, H. *et al.* (2020) 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 306–312. doi: 10.1016/j.oooo.2020.06.021.
50. Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286.
51. Sekar, D. *et al.* (2019) 'Methylation-dependent circulating microRNA 510 in preeclampsia patients', *Hypertension research: official journal of the Japanese Society of Hypertension*, 42(10), pp. 1647–1648.
52. Seppan, P. *et al.* (2018) 'Therapeutic potential of Mucunapruriens (Linn.) on ageing induced damage in dorsal nerve of the penis and its implication on erectile function: an experimental study using albino rats', *The aging male: the official journal of the International Society for the Study of the Aging Male*, pp. 1–14.
53. Sharma, P. *et al.* (2019) 'Emerging trends in the novel drug delivery approaches for the treatment of lung cancer', *Chemico-biological interactions*, 309, p. 108720.
54. Siddique, R. *et al.* (2019) 'Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi', *Journal of conservative dentistry: JCD*, 22(1), pp. 40–47.
55. Somasundaram, S. (2015) 'Fluoride Content of Bottled Drinking Water in Chennai, Tamilnadu', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2015/14691.6594.
56. Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and

- oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306.
57. Subramanyam, D. *et al.*(2018) 'Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries', *European Journal of Dentistry*, pp. 067–070.doi: 10.4103/ejd.ejd\_266\_17.
  58. Teja, K. V., Ramesh, S. and Priya, V. (2018) 'Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study', *Journal of conservative dentistry: JCD*, 21(6), pp. 592–596.
  59. Tulsani, S. G., Chikkanarasaiah, N. and Bethur, S. (2014) 'An in vivo comparison of antimicrobial efficacy of sodium hypochlorite and Biopure MTAD<sup>TM</sup> against enterococcus faecalis in primary teeth: A qPCR study', *The Journal of clinical pediatric dentistry*, 39(1), pp. 30–34.
  60. Türkün, M., Gökay, N. and Özdemir, N. (1998) 'FARKLI ENDODONTİK YIKAMA SOLÜSYONLARININ TOKSİK VE NEKROTİK DOKU ÇÖZÜCÜ ETKİLERİNİN KARŞILAŞTIRMALI OLARAK İNCELENMESİ-COMPARATIVE INVESTIGATION OF THE TOXIC AND NECROTIC TISSUE-DISSOLVING EFFECTS OF DIFFERENT ENDODONTIC IRRIGANTS', *Journal of Istanbul University Faculty of Dentistry*, 32(2), pp. 87–94.
  61. Vasconcelos, B. C., Luna-Cruz, S. M. and De-Deus, G. (2007) 'Cleaning ability of chlorhexidine gel and sodium hypochlorite associated or not with EDTA as root canal irrigants: a scanning electron microscopy study', *Journal of applied oral science: revista FOB*. Available at: [http://www.scielo.br/scielo.php?pid=S1678-77572007000500003&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S1678-77572007000500003&script=sci_arttext).
  62. VijayashreePriyadharsini, J. (2019) 'In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens', *Journal of periodontology*, 90(12), pp. 1441–1448.
  63. VijayashreePriyadharsini, J., SmilineGirija, A. S. and Paramasivam, A. (2018) 'In silico analysis of virulence genes in an emerging dental pathogen A. baumannii and related species', *Archives of oral biology*, 94, pp. 93–98.
  64. Yesilsoy, C. *et al.*(1995) 'Antimicrobial and toxic effects of established and potential root canal irrigants', *Journal of endodontics*, 21(10), pp. 513–515.
  65. Yoshida, T. *et al.*(1995) 'Clinical evaluation of the efficacy of EDTA solution as an endodontic irrigant', *Journal of endodontics*, 21(12), pp. 592–593.



Graph 1: Bar Graph shows association between the type of Irrigant that is used and the age of the patient. X axis represents age of patients and Y axis represents number of teeth that underwent pulpectomy. Most commonly used irrigant in all age groups is Saline. There is a statistical significant association (Chi Square Test, Value=20.898, df=6 and  $p=0.02$ ) which infers that the usage of other irrigants other than saline decreases with age.



Graph 2: Bar Graph shows association between the irrigant that is used for pulpectomy and gender of the patient. X axis represents the gender of the patient and Y axis represents the


number of teeth that underwent pulpectomy. Most commonly used irrigant among both the genders was saline (blue). However there was no statistically significant association between gender of patient and irrigant (Chi Square Test, Value= 5.255, df=3, p=0.154)

	Irrigant				Total
	Saline	Saline, EDTA	EDTA	Saline, Sodium Hypochlorite	
Tooth Number 51	69	44	7	0	120
52	76	48	14	0	138
53	35	14	2	0	51
54	107	66	19	0	192
55	35	25	8	0	68
61	66	47	11	0	124
62	70	44	14	0	128
63	34	18	6	0	58

64	114	74	16	0	204
65	55	31	9	0	95
71	6	1	0	0	7
72	9	1	2	0	12
73	20	9	2	0	31
74	167	84	25	2	278
75	178	88	26	0	292
81	6	2	0	0	8
82	8	4	0	0	12
83	19	12	1	0	32
84	184	89	18	0	291
85	181	92	24	0	297

Total	1439	793	204	2	2438
(p= 0.002)					

Table 1: Table shows association between the teeth number and the Irrigant which is used for pulpectomy. The most commonly used Irrigant is saline. Saline with EDTA and EDTA alone are used more commonly in molars than in anteriors. Saline along with hypochlorite is used only in the lower left molar. (Chi Square Test, Value= 92.578, df=57, p=0.002)



Graph 3: Bar Graph shows association between the level of education and the Irrigant that is used for pulpectomy. X axis represents the level of education and Y axis represents the number of teeth that underwent pulpectomy. The most commonly irrigant used in both levels of education was saline (blue). There is a statistically significant association (Chi Square Test, Value=17.073, df=3 and p=0.001) between the level of education and the Irrigant used for pulpectomy.